

RISK OF DECOMPRESSION SICKNESS IN SHALLOW NO-STOP AIR DIVING:

AN ANALYSIS OF NAVAL SAFETY CENTER DATA 1990-1994

E. T. Flynn E. C. Parker R. Ball

Naval Medical Research and Development Command Bethesda, Maryland 20889-5606

Bureau of Medicine and Surgery Department of the Navy Washington, DC 20372-5120

20000824 145

Approved for public release; distribution is unlimited

NOTICES

The opinions and assertions contained herein are the private ones of the writer and are not to be construed as official or reflecting the views of the naval service at large.

When U. S. Government drawings, specifications, and other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever. The fact that the Government may have formulated, furnished or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Please do not request copies of this report from the Naval Medical Research Institute. Additional copies may be purchased from:

Office of the Under Secretary of Defense (Acquisition & Technology)
Defense Technical Information Center
8725 John J. Kingman Road, Suite 0944
Ft. Belvoir, VA 22060-6218

Federal Government agencies and their contractors registered with the Defense Technical Information Center should direct requests for copies of this report to:

TECHNICAL REVIEW AND APPROVAL

NMRI 98-08

The experiments reported herein were conducted according to the principles set forth in the current edition of the "Guide for the Care and Use of Laboratory Animals," Institute of Laboratory Animals Resources, National Research Council.

This technical report has been reviewed by the NMRI scientific and public affairs staff and is approved for publication. It is releasable to the National Technical Information Service where it will be available to the general public, including foreign nations.

THOMAS J. CONTRERAS CAPT, MSC, USN Commanding Officer Naval Medical Research Institute

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services. Directorate for information Operations and Reports, 1215 Jefferson Daus Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

Latis Highway, suite 1204, Armigion, VA 22202-30			
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE May 1998	3. REPORT TYPE AND TECHNICAL ME	ay 1995 - April 1996
4. TITLE AND SUBTITLE			5. FUNDING NUMBERS
RISK OF DECOMPRESSION S	ICKNESS IN SHALLOW N	O-STOP AIR	PE - 63713N
DIVING: AN ANALYSIS OF	NAVAL SAFETY CENTER	DATA 1990-1994.	PR - MO099
			TAOlA
6. AUTHOR(S)			WU -1510
Flynn, E.T., E.C. Parke	r, and R. Ball		
7. PERFORMING ORGANIZATION NAM	E(S) AND ADDRESS(ES)		8. PERFORMING ORGANIZATION REPORT NUMBER
Naval Medical Research Instit	tute		REPORT NOWIGEN
Commanding Officer 8901 Wisconsin Avenue	•		NMRI 98 08
Bethesda, Maryland 20889-560	17		11WIKI 98- 00
Deffiesda, Maryland 20009-500)/		
9. SPONSORING/MONITORING AGENC	Y NAME(S) AND ADDRESS(ES)		10. SPONSORING / MONITORING
Naval Medical Research and	Development Command		AGENCY REPORT NUMBER
National Naval Medical Cent			DN241126
Building 1, Tower 12			DNZ-1120
8901 Wisconsin Avenue		·	
Bethesda, Maryland 20889-560)6		
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION / AVAILABILITY ST	ATEMENT		12b. DISTRIBUTION CODE
Approved for public release;	distribution is unlimited.		
·			ļ
13. ABSTRACT (Maximum 200 words)			<u> </u>
	SC) dive and incident rep	orts for calendar ve	ars 1990-1994 were
analyzed to determine the r			
using current U.S. Navy no			
performed 163,400 no-deco			_
period and reported 48 incident	-		
cases/1000 dives. There wa			
The DCS rate in the 4th qua		_	
nearly six times the rate in	<u>-</u>		
even in the 4th quartile of U	Jana / no-decompression	i ume was still quit	e low in adsolute terms.
		1 1 11 170	HDGC TILL I
, , ,	-		e II DCS. This is in marked
contrast to previous reports	of military diving operat	nons where Type I	cases predominate.
14. SUBJECT TERMS			15. NUMBER OF PAGES
decompression sickness,	shallow diving, sta	ndard air tables	
USN93 Tables			16. PRICE CODE
	. SECURITY CLASSIFICATION	19. SECURITY CLASSIF OF ABSTRACT	ICATION 20. LIMITATION OF ABSTRAC
OF REPORT Unclassified	OF THIS PAGE Unclassified	Unclassified	Unlimited

The potential impact of adopting new and more restrictive no-decompression limits (USN93) was examined. If the new limits were adopted in the context of single-level dive tables similar to the current Standard Air Tables, and if the pattern of Navy diving remained unchanged, approximately 2.9% of the no-decompression dives conducted between 21 and 55 fsw would be converted to decompression-requiring dives. Approximately 954 dives would be affected annually, adding 419 hours of decompression time to operations each year assuming the decompression was computed using the proposed new USN93 tables. This is an average of 26.3 minutes of decompression for each dive over the new limit. The observed DCS case rate for dives over the proposed USN93 limit is 1.26 cases/1000 dives, a rate very similar to the USN57 4th quartile rate. With the institution of decompression for these dives, the DCS case rate could be expected to fall from 1.26 cases/1000 dives to 1.07 cases/1000 dives, a 15% reduction. In absolute terms, this is an expected reduction of 0.18 case/year.

TABLE OF CONTENTS

	Page
Acknowledgements	iii
Background	1
Methods	3
Results	4
Discussion	18
Impact of Adopting USN93 No-Decompression Limits	19
Conclusions	21
References	22
Tables	
Table 1. Comparison of USN57 and USN93 No-Decompression Limits	2
Table 2. Refinement of the Naval Safety Center Data Set	5
Table 3. Distribution of Dives by Depth Range and Performer	6
Table 4. Air No-Decompression Dives Longer than Six Hours	10
Table 5. Number of Dives Between USN93 and USN57 No-Decompression Limits	10
Table 6. Distribution of Incidents by Bottom Time	12
Table 7. Cases Occurring Beyond the USN93 No-Decompression Limit	14
Table 8. DCS Rate as a Function of Depth Range	14
Table 9. DCS Rate as a Function of Quartile of No-Decompression Time	16
Table 10. DCS Rate for Dives Shorter or Longer than the USN93 Limit	17

Figures

Figure 1.	Frequency Distribution of No-Decompression Dives by Bottom Time	7
Figure 2.	Frequency Distribution of Dives by Maximum Reported Depth	8
Figure 3.	Frequency Distribution of Dives by Last Digit of Reported Bottom Time	9
Figure 4.	Percentage of Cases Observed as a Function of the Fraction of USN57 No-Decompression Time	13
Figure 5.	Cumulative Number of Cases versus Cumulative Number of Dives	17
	Appendices	
Appendix	A Frequency of Shallow Air No-Decompression Diving (1990 - 1994)	24
Appendix	B DCS Case Summary and Descriptions	35

ACKNOWLEDGEMENTS

We greatly appreciate the cooperation of the Naval Safety Center in this endeavor. LCDR Patrick E. Keller USN was instrumental in the organization of the study and facilitated the interaction between our two organizations. Ms. Glenda Batte provided all the diving records, while the staff of the Diving and Salvage Division, especially EMCM(MDV) M. Bonnin USN, spent many hours organizing and verifying the accident reports. Without their help this study would not have been possible.

This work was supported by Naval Medical Research and Development Command Work Unit No. 63713N M0099.01A-1510. The opinions and assertions contained herein are the private ones of the authors and are not to be construed as official or reflecting the views of the Navy Department or the naval service at large.

BACKGROUND

During the 1980s, surveys of operational diving practice (1-4) as well as experimental work at NEDU (5) demonstrated that decompression was inadequate for long, deep air dives. When performed with either in-water decompression or surface decompression techniques, these dives were associated with an unacceptable bends rate (5). Experimentally, Thalmann demonstrated that a 150 feet of sea water (fsw) for 40 min dive or a 190 fsw for 30 min dive required 1.6 times the decompression time called for in the U.S. Naby Standard Air Decompression Table (USN57). Similarly, a 60 fsw/180 min dive required 2.7 times as much decompression (5). In the mid 1980's, concern about the safety of air decompression was so great that the U.K. Department of Energy issued two Diving Safety Memoranda (6) that severely restricted the depth-time envelope for air diving in the North Sea.

In response to these concerns, and to restore the operational envelope for air diving, new decompression models and procedures were developed by several groups including the U.S. Navy, the French Navy, COMEX S.A. (subsequently adopted by the French Ministry of Health), the Canadian Forces, the University of Pennsylvania (for the U.K. Health and Safety Executive and SubSea International, Inc.), and Duke University (for its support of Turkish archaeologic expeditions). Some of these new decompression models continued with the traditional Haldanian (17) approach while others incorporated probabilistic features and risk based on computed bubble dynamics.

The U.S. Navy developed and extensively tested a new probabilistic decompression model in 1993 (7-9). This model, referred to as USN93, produced a new set of air decompression procedures that corrected perceived deficiencies in the decompression from long, deep dives. However, this model also shortened the allowed no-decompression bottom times for shallow dives when compared to USN57. At 40 fsw, for example, USN93 shortened the no-decompression limit from 200 minutes to 144 minutes. At 35 fsw, the no-decompression limit was shortened from 310 to 185 minutes. Table 1 summarizes the changes.

The USN93 model recommendations for shortening the shallow no-decompression limits were consistent with the recommendations of others. The new COMEX tables set a no-

decompression limit of 165 minutes at 40 fsw, while the new University of Pennsylvania tables allowed only 120 minutes. Nevertheless, the need for shorter no-decompression limits was seriously questioned by the Fleet (10). The Fleet maintained that no-decompression diving in the U.S. Navy had an excellent safety record and that shorter no-decompression limits were both mission-compromising and unnecessary.

As a part of its review of USN93-based decompression procedures for Fleet introduction, the Naval Sea Systems Command (NAVSEA) (11) requested that the Naval Medical Research Institute (NMRI) assess the historical DCS rate for shallow, nodecompression air diving in the U.S. Navy using the USN57 procedures. This paper reports the results of that assessment and estimates the potential impact of adopting the USN93 nodecompression limits.

Table 1
Comparison of USN57 and USN93 No-Decompression Limits for Shallow Water Air Diving

Depth Range (fsw)	USN57 (min)	USN93 (min)
21-25	Unlimited	338
26-30	Unlimited	245
31-35	310	185
36-40	200	144
41-45	100	114
46-50	100	93
51-55	60	77
56-60	60	64

METHODS

The assessment was performed using data archived in the Naval Safety Center (NSC). The NSC was asked to identify all air dives in its database that met the following criteria:

- a. Were performed in the five-year period between 01 January 1990 and 31 December 1994.
- b. Were performed by Navy, Marine, Reserve, or Civilian divers as coded in Box A-10 of DD Form 2544.
- c. Were performed in open water as coded in Box B-16.
- d. Were to a maximum depth of 21-55 fsw as coded in Box B-29.
- e. Were labelled as "Air Diving, No-decompression", code "A" in Box C-34.

Dives reported to NSC by other Department of Defense components (e.g. Army, Coast Guard) were not included because incident reporting for these components was felt to be incomplete (M. Bonnin, NSC; personal communication).

Once received from NSC, the data set was scanned for possible errors including duplicate entries, dives with a breathing apparatus not compatible with air diving (LAR V, Mark 15, Mark 16), dives not on 21% oxygen or showing helium in Box B-26, and dives with bottom times exceeding the air no-decompression limits. Dives failing to pass these additional screens were deleted and the remaining dives were aggregated into a spreadsheet in depth increments of 5 fsw from 21-55 fsw and in time increments of both 1 and 10 minutes from 0-360 min. For dives in the 21-25 fsw and 26-30 fsw depth ranges, additional time categories of 6-8 h, 8-10 h, 10-12 h and >12 h were added to accommodate dives longer than 360 min. From this spreadsheet, frequency distributions of dives by depth, bottom time and performer were created.

The NSC was also asked to provide copies of all incident reports bearing a diagnosis of either decompression sickness or arterial gas embolism for the period 01 January 1990 to 31 December 1994. These reports were examined to select those incidents that occurred on open-water, air, no-decompression dives between 21 and 55 fsw. These incidents were then reviewed by a panel of five NMRI undersea medical officers who segregated them by consensus vote into one of three categories: (1) Definitely Decompression Illness (DCI), (2) Definitely not DCI, and (3) Uncertain whether it is DCI or not. Category 1, Definitely DCI, was further divided into three subclasses: (a) Decompression Sickness (DCS), (b) Arterial Gas

Embolism (AGE), and (c) DCS or AGE. This categorization scheme recognizes the fact that it is frequently impossible to tell whether an illness observed following decompression represents DCS or AGE (12). The selected incidents were also checked against the dive database to ascertain that the causative dive was included in the database and the depths and bottom times of the dive and the accident reports matched.

Prior to any analyses, all incidents categorized as Definitely not DCI by the panel of undersea medical officers were eliminated from the data set. Also, those incidents where both NSC and the panel of undersea medical officers agreed that the case was a definite AGE were eliminated. This left only those cases where the diagnosis was either (1) Definite Decompression Sickness or (2) Possible Decompression Sickness. All equivocal cases that might have been decompression sickness were thus retained in the analysis.

The DCS rate was computed by dividing the number of cases observed by the number of dives performed, then multiplying the quotient by 1000 to express the rate in cases/1000 dives. The DCS rate was computed by depth band alone (e.g., 21-25 fsw), by depth band and whether the dive was shorter or longer than the USN93 no-decompression limit, and by depth band and whether this dive was in the first, second, third, or fourth quartile of the USN57 no-decompression time. Rates across depth bands were also computed. The relative risk of one condition compared with another was computed by dividing the rate in that condition by the rate in the reference condition. Tests of statistical significance were not conducted because of the small number of DCS cases in the data set.

RESULTS

The NSC identified 166,443 dives based on the criteria (a - e) outlined above. Of these dives, 863 were found to be duplicate entries, 223 to have bottom times exceeding the current no-decompression limit, and 1,957 that were not entirely air dives. These invalid entries represented approximately 2% of the data received from NSC and when they were eliminated from the data set, 163,400 dives remained for analysis. The refinement of the NSC data set is summarized in Table 2.

Table 2
Refinement of the Naval Safety Center Data set

Initial NSC Total	166,443	
Eliminate Non-Air Dives		
Mark 15	154 414 287 90 12	-1,957
Eliminate Dives Longer than No-D Li	mit	-223
Eliminate Duplicate Entries		-863
Final NSC Total		163,400

Duplicate entries represented one-half of one percent of the data set. Usually two entries were made, one with the diver's full name and one with his initials. In one instance, two dives with the same diver had an identical date and surfacing time, but the time recorded for leaving the surface differed by 2 minutes.

A number of dives (n=223; 31-55 fsw) listed as "no-decompression" actually exceeded the USN57 no-decompression limit. In some instances, the reported bottom time was just over the limit. Some reported dives, however, were well over the limit. At 36-40 fsw, for example, 16 dives were reported with bottom times of 271-290 minutes (No-decompression limit = 200 min). Across the depth range of 31-55 fsw, 28 "no-decompression" dives were reported with bottom times in excess of 6 hours!

A unique situation was discovered in 32 records. In these records, the day of the month the dive was performed was entered into the day column of the bottom time field, resulting, for example, in a no-decompression dive to 32 fsw for 23 days, 0 hours, and 13 minutes. Such errors were corrected and these dives were retained in the database. They are not part of the 223 dives over the limit described above.

Dives performed with the LAR V, Mark 15, and Mark 16 and listed as air no-decompression may have been SPECWAR dives in which these rigs were used for part of the time in what otherwise was an air dive. Alternatively, they may have been no-decompression dives no those respective rigs in which code "A" in Box C-34 of DD 2544 was checked in error. No attempt was made to verify the nature of these dives. They were eliminated from the dataset.

Table 3 shows the distribution of the 163,400 dives by depth range and performer. More detailed information is given in Appendix A. The greatest number of dives occurred in the 26-30 fsw depth range. Eighty-two percent of the dives were between 21 and 40 fsw. Active duty and civilian Navy divers performed more than 90% of the dives. Within Navy, the ratio of active duty dives to civilian dives was approximately 3 to 1.

Table 3

Distribution of Dives by Depth Range and Performer

Depth Range	Total Dives	Navy	Civilian	Marine	Reserve
(fsw)			(% within d	lepth range)	
21-25	29,325	75.8	13.4	4.7	6.2
26-30	43,455	68.0	24.7	2.4	4.9
31-35	28,621	72.4	22.2	1.8	3.5
36-40	33,198	64.1	29.8	1.7	4.4
41-45	11,415	75.6	18.0	1.8	4.6
46-50	13,302	65.2	28.5	2.0	4.2
51-55	4,084	78.5	12.2	3.6	5.7
Total	163,400	70.0	22.8	2.5	4.7

Figure 1 shows the frequency distribution of dives across all performers as a function of bottom time. For all depth ranges under study, the frequency rises rapidly with increasing bottom time, reaching a peak at 21-30 minutes. Frequency then declines with a long tail until the no-decompression time is reached. A secondary peak in dive frequency is not seen as the no-decompression limit is approached. This suggests that divers are not pushing the no-decompression limit. Appendix A gives a further breakdown of these time distributions by category of performer.




Figure 1. Frequency Distribution of No-Decompression Dives as a Function of Bottom Time. The number of dives performed in each 10 minute increment of bottom time from zero to the USN57 limit is shown for all 7 depth ranges studied. Numbers are the sum of values for Navy, Civilian, Marine, and Reserve divers. At all depths the highest frequency is observed between 21 and 30 min of bottom time.

Figure 2 shows the frequency distribution of dives by the maximum reported depth (Box B-29, DD 2544). Substantial spikes in dive frequency are observed at increments of 5 and 10 fsw, suggesting rounding up of the actual maximum depth of the dive to the next deeper 5- or 10-fsw depth. Spikes at the 10-fsw mark consistently exceed those at the 5-fsw mark, suggesting further that rounding up to the next deeper 10-fsw depth is the more common practice.




Figure 2. Frequency Distribution of Dives by Maximum Reported Depth. Pronounced peaks are observed at 5- and 10-fsw intervals indicating rounding up of maximum depth to the next 5- or 10-fsw mark.

Figure 3 shows the frequency distribution of dives by the last digit of the reported bottom time (Box B-30, DD 2544). Zero represents dives of 10, 20, 30 min, etc, while 5 represents dives of 5, 15, 25 min, etc. A substantial spike is present at 0, suggesting rounding up the bottom time to the next 10-minute mark. There also appears to be some rounding up to the next 5 minutes as well, although the evidence for this is less strong.




Figure 3. Frequency Distribution of Dives by Last Digit of Reported Bottom Time. The highest frequencies are observed at 5 and 0 suggesting some rounding up of bottom time to the next 5 or 10 minute mark.

For dives to 30 fsw and shallower, the USN57 no-decompression limit is infinite. Table 4 shows the number of dives between 21and 30 fsw that were reported to be greater than 6-hours duration. Fifty-eight dives were identified in this category; 11 of these dives had reported durations greater than 12 hours.

Table 5 shows the number of dives reported between the USN93 no-decompression

limit and the USN57 limit for the depth range indicated. The blank entries for 41-45 and 51-55 fsw reflect the fact that the USN93 limit is more permissive than the USN57 limit in these ranges. All dives conducted between the USN93 limit and the USN57 limits in these two ranges therefore would have been labelled decompression dives under current rules and as such were not captured for this analysis.

Table 4
Air No-Decompression Dives Longer than Six Hours

Depth Range (fsw)	Duration (hours)				Total
	6-8	8-10	10-12	>12	
21-25	16	3	2	5	26
26-30	13*	7	6	6	32
Total	29	10	8	11	58

^{*}Two incidents reported (see Table 7 below)

Table 5
Number of Dives Between USN93 and USN57 No-Decompression Limits

Depth Range (fsw)	≤ USN93	≤USN57	Between
21-25	29,291	29,325	34
26-30	42,791	43,455	664
31-35	26,970	28,621	1651
36-40	31,143	33,198	2055
41-45			
46-50	12,976	13,302	366
51-55			
Total			4770

Approximately 4,770 dives between 21 and 55 fsw would be converted from nodecompression dives to decompression dives in a 5 year period by adopting the USN93 limit in a single level dive format (Table 5). This represents 2.9% of the total dives conducted in this depth range. The word "approximately" is used because the dives falling between the USN93 and USN57 limits in the depth ranges of 41-45 and 51-55 fsw would actually be converted in the opposite direction, from decompression dives to no-decompression dives, making the total number of dives converted to decompression dives over the full depth range of 21-55 fsw less than 4,770. The precise reduction from 4,770 cannot be determined.

A total of 60 DCS/AGE incidents on open-water, air no-decompression dives were identified from the material provided by NSC. Of these, NSC considered 31 cases to represent DCS while 29 were considered AGE. Of the 31 DCS cases, 27 (87%) were reported by NSC as Type II DCS.

The NMRI panel of undersea medical officers felt that 36 of the 60 cases were definitely decompression-related (Definitely DCI), 7 were definitely not decompression-related (Definitely not DCI), and 17 were equivocal (*i.e.*, they may or may not have been decompression-related). For the cases labelled Definitely not DCI, symptoms were already present during the bottom phase of the dive or the diver failed to show any response to recompression therapy. Of those cases labelled Definitely DCI by the panel, 18 were felt to be DCS, 5 to be AGE, and 13 to be either DCS or AGE. After eliminating 7 cases considered Definitely Not DCI and 5 cases where both NMRI and NSC agreed on the diagnosis of AGE, 48 cases remained for analysis. One of the 7 cases eliminated as definitely not DCI had a bottom time longer than the USN93 limit (265 min at 28 fsw). The remaining 6 definitely not DCI cases and the 5 AGE cases all had bottom times shorter than the USN93 limit.

There was generally very good agreement between the depth/bottom time information provided on the incident report and the information contained in the dive database for the causative dive. There were some instances, however, where there were discrepancies. In one instance, the dive was reported as 90 minutes; the incident report indicated 91 minutes. In another instance, the dive record and the incident report numerical data both indicated a depth of 22 fsw, but the incident report narrative indicated a depth of 29 fsw. In a third instance, the dive record and incident report numerical data indicated a bottom time of 52 minutes, but the incident report narrative indicated 51 minutes. In a fourth instance, the dive record and

incident report numerical data indicated a depth of 26 fsw, while the incident report narrative indicated a depth of 30 fsw. Since there was agreement between the dive record and the incident report numerical data in the latter three instances, these values were used in the analysis. In the first instance, the 90-minute bottom time from the dive record was used.

Table 6 shows the distribution of incidents by bottom time within each depth range. Cases indicated by bold print lie beyond the USN93 limit. The shortest bottom time for a case was 13 minutes, the longest was 397 minutes. Within each depth range, cases appeared across the range of bottom times, with no evidence of clustering near the no-decompression limits.

Table 6
Distribution of Incidents by Bottom Time

Depth Range (fsw)	Bottom Time (min)
21-25	29,40,40,44,44,63,90,140
26-30	17,22,25,30,45,50,61,73,74,135,397,397
31-35	43,52,52,76,115,143,191,278
36-40	20,30,50,74,91,113,121,134,176,189
41-45	15,46,97
46-50	40,52,52,57
51-55	13,46,58

Figure 4 shows the percentage of total cases observed in the 26-30, 31-35, and 36-40 fsw depth ranges as a function of the bottom time expressed as percentage of the USN57 nodecompression limit. These are the depth ranges where implementation of the USN93 limit would have the greatest impact (see Table 5). For purposes of this analysis the USN57 nodecompression limit at 26-30 fsw was considered to be 400 minutes; in fact, it is infinite. For all three depth ranges, 70% or more of the cases are observed in the first 60% of the nodecompression time. For the 26-30 fsw range, nearly 80% of the cases are observed in the first 20% of the no-decompression time. (Note: This result would be even more striking if a nodecompression limit greater than 400 min had been selected for this depth range.) The

unexpectedly large number of cases at bottom times well short of the no-decompression limit is a reflection of the very large number of dives performed at these short times (see also the discussion below).

Six of the 48 cases (12.5%) occurred at bottom times between the USN93 limit and the USN57 limit. Table 7 summarizes the essential features of these cases. Appendix B gives greater detail. Three of the cases were from ship husbandry operations; three from special-warfare operations. All but one case demonstrated some form of neurological involvement. The NSC labelled two of the cases AGE, the remaining four DCS. The NMRI review panel felt one of the cases labelled AGE by NSC was in fact DCS; the other case labelled AGE by NSC was considered equivocal (*i.e.*, could have represented either DCS, AGE, or was not-decompression related). The first two cases occurred on the same dive.




Figure 4. Percentage of Cases Observed as a Function of the Fraction of USN57 No-Decompression Time. Most of the reported cases are observed well before the no-decompression limit is neared.

Table 7
Cases Occurring Beyond the USN93 No-Decompression Limit

Dive (fsw/min)	Purpose	Symptoms	Diagnosis	
			NSC	NMRI
30/397	SW/SDV	Neuropsychiatric	DCS TYPE 2	EQV
30/397	SW/SDV	Numbness	DCS TYPE 2	DCS
33/278	SW/DDS	Back Pain	DCS TYPE 1	DCS
34/191	SH	Hearing Loss	AGE	EQV
37/176	SH	Pain/Weakness	AGE	DCS
40/189	SH	Pain/Numbness	DCS TYPE 2	DCS

SW-special warfare; SDV-swimmer diver vehicle; DDS-dry deck shelter; EQV-equivocal case, may or may not have been decompression-related.

Table 8 shows the DCS rate (cases/1000 dives) by depth interval without regard to bottom time. The rate is remarkably constant, ranging between 0.26 and 0.30 cases/1000 dives (2.6-3.0 cases/10,000 dives) across the entire range of 21 to 50 fsw. Only between 51 and 55 fsw does the rate increase to 0.73 cases/1000 dives.

Table 8
DCS Rate as a Function of Depth Range

Depth Range (fsw)	Bottom Time (min)	Total Dives (#)	Total Cases (#)	Rate (cases/1000)
21-25	0-INF	29,325	8	0.27
26-30	0-INF	43,455	12	0.28
31-35	0-310	28,621	8	0.28
36-40	0-200	33,198	10	0.30
41-45	0-100	11,415	3	0.26
46-50	0-100	13,302	4	0.30
51-55	0-60	4,084	3	0.73
Total		163,400	48	0.29

Table 9 shows the DCS rate further segregated into quartiles of USN57 no-decompression time. At 21-30 fsw, where the USN57 limit is infinite, four quartiles of 100 minutes each were arbitrarily chosen, allowing the analysis to cover 400 minutes of bottom time and to encompass all of the observed cases of DCS. This process excluded 43 dives beyond the 400-minute mark from the analysis.

Overall, the quartile analysis shows a steadily increasing DCS rate from quartile to quartile, ranging from 0.22 cases/1000 dives in the first quartile to 1.28 cases/1000 dives in the fourth quartile. The DCS rate in the fourth quartile is 2.2 times the rate in the 3rd quartile and 5.8 times the rate in the first quartile.

Figure 5 plots the cumulative number of DCS cases at 26-30, 31-35, and 36-40 fsw as a function of the cumulative number of dives undertaken beginning at a bottom time of 1-10 minutes and then proceeding to progressively longer bottom times. If the DCS rate was a constant value independent of bottom time, these plots would be expected to be straight lines with a y-intercept of zero and a slope equal to the DCS rate. At each depth, the curves bow upward indicating a rising DCS rate with increasing bottom time. The information plotted in Figure 5 is from Appendix A and Table 6.

Table 10 shows the DCS rate within each depth range when the dives are segregated into those equal to or shorter than the USN93 limit and those longer than the USN93 limit. The latter are the dives that would be converted to decompression dives by adopting the USN93 limit. The depth ranges 41-45 and 51-55 fsw are not shown since the USN93 limit exceeds the USN57 limit in these ranges. For dives equal to or shorter than the USN93 limit, the mean DCS rate is 0.26 cases/1000 dives (range 0.22-0.31). Longer than the USN93 limit, the mean DCS rate is 1.26 cases/1000 dives (range 0-3.01), a 4.8-fold increase over those dives equal to or shorter than the limit.

Table 9

DCS Rate as a Function of Quartile of No-Decompression Time

Depth Range	Bottom Time	Total Dives	Total Cases	Rate
(fsw)	(min)	(#)	(#)	(#/1000)
<u></u>	0-100	24,584	7	0.28
21-25	101-200	3,867	1	0.26
21 23	201-300	775	0	0.00
	301-400	80	0	0.00
**************************************	0-100	36,641	9	0.25
26-30	101-200	5,370	1	0.19
20-30	201-300	1,330	0	0.00
	301-400	90	2	22.22
	0-78	20,701	4	0.19
31-35	79-155	5,175	2	0.39
31-33	156-233	1,981	1	0.50
	234-310	764	1	1.31
	0-50	21,300	3	0.14
36-40	51-100	7,374	2	0.27
30 10	101-150	2,704	3	1.11
	151-200	1,820	2	1.10
	0-25	3,826	1	0.26
41-45	26-50	4,646	1	0.22
11 15	51-75	1,743	0	0.00
·	76-100	1,200	1	0.83
	0-25	3,862	0	0.00
46-50	26-50	5,679	1	0.18
10.50	51-75	2,347	3	1.28
	76-100	1,414	0	0.00
	0-15	651	1	1.54
51-55	16-30	1,394	0	0.00
31-33	31-45	1,139	0	0.00
	46-60	900	2	2.22
	1 st Quartile	111,565	25	0.22
Summary	2 nd Quartile	33,505	8	0.24
	3 rd Quartile	12,019	7	0.58
	4 th Quartile	6,268	8	1.28
TOTAL		163,357	48	0.29




Figure 5. Cumulative Number of Cases versus Cumulative Number of Dives Undertaken. At the three depths shown, the curves tend to bow upward suggesting the DCS rate is rising with increasing bottom time.

Table 10
DCS Rate for Dives Shorter or Longer than the USN93 Limit

Depth Range (fsw)	Shorter	than USN9	3 Limit	Longer than USN93 Limit			
	Dives (#)	(#) Cases (#) Rate (#/1000)		Dives (#)	Cases (#)	Rate (#/1000)	
21-25	29,291	8	0.27	34	0	0.00	
26-30	42,791	10	0.23	664	2	3.01	
31-35	26,970	6	0.22	1651	2	1.21	
36-40	31,143	8	0.26	2055	2	0.97	
41-45	11,415	3	0.26		-		
46-50	12,936	4	0.31	366	0	0.00	
51-55	4,084	3	0.73		-		
Total	158,630	42	0.26	4770	6	1.26	

DISCUSSION

The epidemiologic data that the NSC collects on U.S. military diving operations is among the best in the world today. Nevertheless, we found that these data sets were not error-free. Approximately 2% of the diving records we examined had to be excluded from the study because they represented duplicate entries, had listed bottom times in excess of the air no-decompression limits, or listed a breathing apparatus inconsistent with air diving (e.g., LAR V). We observed that reported maximum depth was usually rounded up to the next even increment of 5 or 10 fsw. Reported bottom time appeared to be rounded up as well.

Incident reports occasionally contained conflicting depth and time information. By far the more limiting feature of the incident report, however, was the quantity and quality of the information contained in the narrative section. These narratives, containing 100-200 words, provided scant information on the clinical presentation of the cases (see Appendix B for examples). The information was so brief that the panel of undersea medical officers in seven cases could not conclude whether the case was related to decompression or not. The narratives appeared to be written largely by non-medical personnel, and in the cases where two divers from the same dive were treated, the narratives were nearly identical for both divers (see Appendix B). A different categorization may have been reached by the panel if full narrative summaries of the incidents written by the attending physician or corpsman had been available for examination.

Eighty-seven percent of the cases reported to NSC as DCS were labelled Type II. The prominence of Type II symptoms was surprising given previous reports of the experimental and operational experience of professional divers (13,14,15) in which Type I symptoms are the dominant manifestation of DCS. The finding of a high prevalence of Type II symptoms is consistent, however, with recently reported observations in recreational divers (15,16) for whom, as a class, medical monitoring and incident reporting are highly variable.

For all depth ranges, the incidents of definite and possible DCS were distributed broadly over the range of bottom times up to the USN57 no-decompression limit. The cases were not clustered near the no-decompression limit as might be expected. In fact, most cases were observed in the first quartile of the no-decompression time, and can be attributed to the

high frequency of diving at these short bottom times (see Figure 1 and Table 9), not to a higher risk level. Experience has shown that DCS can occur after almost any hyperbaric exposure, even very short and shallow ones. Although the probability of DCS on these dives is very low, the sheer number of dives makes the observance of some cases inevitable.

The data show a clear pattern of increasing risk of DCS with increasing bottom time. The DCS rate for the 4th quartile of no-decompression time is 2.2 times the 3rd quartile rate and nearly 6 times the 1st quartile rate. The DCS rate for dives longer than the USN93 limit is 4.8 times the rate for dives shorter than the limit.

Despite the pattern of increasing risk with increasing bottom time, the absolute risk of DCS during no-decompression diving as currently practiced is quite low. For dives falling between the USN93 limit and the USN57 limit, the mean DCS rate is only 1.26 cases/1000 dives. The highest rate observed in this interval is 3.01 cases/1000 dives.

It would have been highly desirable to make point estimates of the DCS rate <u>at</u> the USN93 and USN57 limit times for each depth range rather than estimates for the preceding interval. The DCS rate for dives conducted <u>at</u> the longer USN57 limit, for example, will be clearly greater than the rate for all the dives between the USN93 and the USN57 limit. Unfortunately, the low number of reported cases would not sustain such an analysis.

IMPACT OF ADOPTING THE USN93 NO-DECOMPRESSION LIMITS

The foregoing analysis allows assessment of the likely operational impact of adopting the USN93 no-decompression limits. If the new limits were adopted in the context of single-level dive tables similar to USN57 and the pattern of diving in the Fleet remained unchanged, approximately 954 dives per year (4,770 dives/5 years, see Table 5) would be converted from no-decompression dives to dives requiring decompression. This number is a slight overestimate since a small number of dives in the 41-45 and 51-55 fsw range now requiring decompression would be converted to no-decompression dives. The 954 dives would add approximately 419 hours of decompression time to operations annually assuming that these dives were all square dives to the maximum depth and bottom time indicated in the NSC data and that decompression was computed using the new USN93 tables. This is an average of 26.3 minutes of decompression per dive for this subset of dives.

The institution of decompression for the 954 dives between the USN93 and USN57 limits would reduce the DCS risk below the current risk with no-decompression, but would not eliminate risk altogether. The decompression tables themselves are associated with a finite level of risk. Using the USN93 algorithm, the risk of performing these 954 dives with and without decompression was computed. For this calculation, the 4,770 dives over the USN93 limit for the five-year period were aggregated into a spreadsheet in depth intervals of 5 fsw (e.g., 26 to 30 fsw) and time intervals of 10 minutes (e.g., 251 to 260 minutes). For all dives in the spreadsheet cell, the depth for the calculation was taken as the upper limit of the cell (e.g., 30 fsw for 26-30 fsw) and the bottom time was taken as the mid point of the cell (e.g., 255 min for 251-260 min). For the 4,770 dives in question, the DCS risk with decompression was estimated by the algorithm to be 85% of the risk of performing them without decompression. In other words, the DCS risk is approximately 15% lower with USN93 decompression than without.

Six cases of DCS were observed in 5 years on the 4770 dives performed over the USN93 limit. This gives a baseline risk of 1.26 cases/1000 dives when these dives are performed without decompression. Based on the analysis outlined above, the risk on these dives with the institution of decompression would be expected to fall to 1.07 cases/1000 dives (1.26 x 0.85). For the 954 dives over the USN93 limit annually, this would result in a rate of 1.02 cases/year with decompression vice 1.20 cases/year without decompression, or the avoidance of 0.18 case/year. This number is a slight overestimate of the benefit since the dives in the 41-45 fsw and 51-55 fsw ranges converted from decompression dives to nodecompression dives would add back in a small risk.

Thus, the best estimate of the cost/benefit of adopting the USN93 limits in the context of single level dive tables is the addition of approximately 419 hours of decompression time annually to operations for the avoidance of 0.18 cases of DCS.

CONCLUSIONS

- 1. The overall DCS rate for shallow water, air, no-stop diving in the U.S. Navy is very low, approximately 0.29 cases/1000 dives. This overall rate remains nearly constant over the depth range of 21-50 fsw.
- 2. The observed probability of DCS with shallow no-decompression diving increases with increasing bottom time. The DCS rate during the 4th quartile of USN57 no-decompression time (1.28 cases/1000 dives) is approximately six times the rate in the 1st quartile (0.22 cases/1000 dives). For bottom times longer than the proposed USN93 limit, the DCS rate is 1.26 cases/1000 dives, which is substantially greater than the DCS rate of 0.26 cases/1000 dives for dives shorter than the limit, but still quite low in absolute terms.
- 3. There is no evidence that divers are currently pushing the no-decompression limits in shallow water. In fact, the vast majority of dives are conducted at less than 25% of the current no-decompression limits for the depth ranges studied here.
- 4. The proportion of DCS cases reported as Type II in shallow no-stop diving is significantly higher than has been previously reported in military divers.
- 5. If the USN93 limits were adopted and used in the context of single-level diving tables, approximately 0.18 cases of DCS would be avoided per year at a cost of approximately 419 hours of additional decompression time.
- 6. These data do not support shortening the USN57 no-decompression limits in shallow water.

REFERENCES

- 1. Berghage, T.E. and Durman, D. <u>U.S. Navy air decompression schedule risk analysis</u>. NMRI #80-1, Naval Medical Research Institute, Bethesda, MD. January 1980.
- 2. Shields, T.G. and Lee, W.B. <u>The incidence of decompression sickness arising from commercial offshore air-diving operations in the U.K. sector of the North Sea during 1982/83.</u> Robert Gordon's Institute of Technology, Abderdeen, Scotland, 1986.
- 3. Shields, T.G., Duff, P.M. and Wilcock, S.E. <u>Decompression sickness from commercial offshore air-diving operations on the U.K. continental shelf during 1982 to 1988</u>. Robert Gordon's Institute of Technology, Aberdeen, Scotland, 1989.
- 4. Imbert, J.P. and Bontoux, M. "Production of procedures: COMEX." In: Nashimoto, I. and Lanphier, E.H., eds. <u>36th Workshop of the Undersea and Hyperbaric Medical Society</u>, Bethesda, MD, 1987.
- 5. Thalmann, E.D. <u>Air-N₂O₂ decompression computer algorithm development</u>. NEDU #8-85, Naval Experimental Diving Unit, Panama City, FL, 1985.
- 6. U.K. Department of Energy, Diving Safety Memorandum 7, 1986 and Diving Safety Memorandum 5, 1988.
- 7. Parker, E.C., Survanshi, S.S., Weathersby, P.K., and Thalmann, E.D. <u>Statistically based decompression tables. VIII. Linear-exponential kinetics</u>. NMRI #92-73, Naval Medical Research Institute, Bethesda, MD, 1992.
- 8. Survanshi, S.S., Weathersby, P.K., and Thalmann, E.D. <u>Statistically based decompression tables X: Real-time decompression algorithm using a probabilistic model</u>. NMRI #96-06, Naval Medical Research Institute, Bethesda, MD, March 1996.
- 9. Thalmann, E.D., Kelleher, P.C., Survanshi, S.S., Parker, E.C., and Weathersby, P.K. Statistically based decompression tables XI: Manned validation of the LE probabilisite decompression model for air and nitrogen-oxygen diving. Joint Naval Medical Research Institute, Bethesda, MD and Naval Experimental Diving Unit, Panama City, FL Technical Report, In preparation.
- 10. Working Divers Conference 95 Proceedings. Naval Sea Systems Command, Washington,

D.C., 28 July 1995.

- 11. NAVSEA ltr 10560 Ser 00C3B/3336 of 19 Sept 95
- 12. Francis, T.J.R. and Smith, D.J., eds. "Describing Decompression Illness." <u>42nd</u> Workshop of the Undersea and Hyperbaric Medical Society, Undersea and Hyperbaric Medical Society, Bethesda, MD, 1991.
- 13. Van der Aue, O.E., Kellar, R.J., and Brinton, E.S. <u>The effect of exercise during decompression from increased barometric pressures on the incidence of decompression sickness in man.</u> NEDU #8-49, Navy Experimental Diving Unit, Washington, D.C., 1949.
- 14. Rivera, J.C. "Decompression sickness among divers: an analysis of 935 cases." Military Medicine, Vol. 129 pp. 314-334, 1964.
- 15. Elliott, D.H. and Moon, R.E. "Manifestations of the decompression disorders." In: Bennett, P.B. and Elliott, D.H., eds., <u>The Physiology and Medicine of Diving</u>. 4th Edition. W.B. Saunders Co., Philadelphia, 1993.
- 16. Arness, M.K., "SCUBA decompression illness and diving fatalities in an overseas military community". <u>Aviation Space and Environmental Medicine</u>, Vol. 68 pp. 325-333, 1997.
- 17. Boycott, A.E., Damant, G.C.C., Haldane, J.S. "The Prevention of Compressed Air Illness". <u>Journal of Hygiene</u> (Camb.) Vol. 8 pp. 342-443, 1908.

APPENDIX A

FREQUENCY OF SHALLOW AIR NO-DECOMPRESSION DIVING

1990 - 1994

Table A1

Distribution of Dives by Depth and Bottom Time

All Dives

	P Beyo	2 # # # # # # # # # # #	113		# Beyo	-282	. 2		# Bayo	2.3	2 - =	2	
	Tobale	16228 21463 13031 16381 7241 8361 2968	ž.		, 1	13071 21960 15610 16875 4232 3962	7		300	28288 43423 28441 33256	11473 13343 4102	163537	
	351-360	n n 0 0 0 0	ω	16228 21463 13031 16361 7241 9361 2968	351-360	n-10000	, o	13071 21960 15610 14875 4272 3962 1134	351-360	6 4 N O	•••	5	28299 43423 28641 33256 11473 113343 4102
	341-350	00000	8	16225 21460 13031 16381 7241 2961 2968	341-350	0-0000		13088 21959 15805 14875 4232 3982 1134	341:350	0 4 0	•••	6	29293 43419 29636 33256 11473 113343 4102
	351-340	00000	51	16225 21459 13030 18381 7241 9361 2968	331-340	00-000	-	13068 21958 15605 14875 4272 3962 1134	331:340	N 0 10 -	0 10 0	13	29293 43417 28635 33256 11473 13343 4102
	321-330		ō	16223 21459 13023 16380 7241 2959 2968	321-330	-00-000		13068 21958 15604 14875 4232 3962 1134	0221-330	4 6 0 -		8	29291 43417 28627 33255 11473 13341 4102
	311-320	B FO 0 0 0 0	5	16220 21450 13017 18380 7241 2958	311-320	200000		13067 21958 15804 14874 4272 3982 1134	311-320	2000	000	g	29287 43408 28621 33254 11473 13341 4102
	301-310	N B B O N O O	5	16212 21447 13017 16378 7741 9359 2968	901-310	* 3 2 0000	145	13051 21955 15604 14074 4232 3962 1134	301-310	6 2 30	000	3	29263 43402 28621 33252 11473 13341 4102
	291-300	4520000	*	16210 21438 13011 18378 7239 8059 2968	291-300	8 5 B 0 0 - 0	217	13016 21903 15546 14874 4232 3962 1134	281-300	75 1.50 0	0-0	255	29226 43341 26657 33252 11471 13341 4102
	281-290	ŭ r ū ◆ 0 0 0	37	16206 21428 12987 16378 7239 9359 2968	281-280	2520000	175	12963 21602 15464 14874 4232 3981 1134	281-290	185	000	212	29189 43230 28451 33252 11471 13340 4102
	271-280	m # 12 # 0 0 0	6	16183 21421 12974 18374 7239 9359 2968	271-280	88200-0	ã	12952 21729 15396 14871 4232 3981 1134	271-280	3 % 38 0	0-0	211	29145 43150 28370 33245 11471 13340 4102
	261-270	20000	18	16185 21413 12947 18368 7239 8359 2968	281-270	422-000	500	12912 21661 15343 14871 4232 3960 1134	281-270	8 8 8 6	000	7	29097 43074 26290 33239 11471 13339
	251-280	27.0000	5	16170 21394 12928 18368 77239 2959	251-280	2880000	82	12868 21580 15260 14870 4232 3980 1134	251-260	\$ £ 50 °	000	308	29038 42974 28188 33236 11471 13339 4102
	241-250	* \tau \tau \tau \tau \tau \tau \tau \tau	8	16152 21357 12907 18366 77239 9359 2968	241-250	\$ \$ \$ 0000	002	12823 21484 15171 14670 4232 3980 1134	241:250	11.4	000	282	28975 42841 28078 33236 11471 13339 4102
	231-240	335 0000	921	16147 21342 12861 18380 7239 9358 2968	231-240	25 E + 0 + 0	2	12777 21365 15086 14870 4232 3980 1134	231-240	817. 84. 84.	0-0	453	28924 42727 27967 33230 11471 13339 4102
	221-230	5854000	2	16113 21268 12644 18355 7239 2958 2968	221-230	3 - 8	275	12695 21260 14875 14866 4232 3979 1134	ZI-230	79 <u>15 00</u> 54	000	336	28808 42548 27819 33221 11471 13338 4102
	211-220	889	87	16094 21258 12634 12634 7239 7239 2968	211-220	5 8 8 v o o o	979	12627 21149 14879 14866 4232 3978 1134	211-220	128 185 25	000	457	28721 42407 27713 33219 11471 13338 4102
	201-210	***	28	16072 21229 12815 18345 17239 9359 2968	201-210	211 25 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	38	12525 21013 14743 14861 4232 3979 1134	201-210	231		233	28597 42242 27558 33206 11471 13338 4102
	191-200	÷28.89000	<u>\$</u>	16038 21153 12785 12785 7239 8459 2968	191-200	25.55.55.000	817	12413 20858 14625 14856 4232 3979 1134	002-161	175 284 285	000	1011	28451 42011 27410 33198 11471 13338 4102
	181-190	*888000	175	15991 21101 12750 16262 7239 9359 2968	181:190	37,82,82,000	30	12285 20653 14376 14621 4232 3979 1134	81-181	208 328 410	000	1239	28276 41754 27126 32903 11471 13338 4102
	171-180	8 2 3 8 0	335	15943 21041 12717 16246 7239 9359 2968	171-180	355 235 0 0	1182	12125 20385 14116 14245 4232 3979 1134		278 418 398 413	n - 0	1517	28068 41426 28833 32483 11471 13338 4102
	161-170	2 2 2 0 0 0	88	15863 20905 12674 18179 7233 9458 2968	071-181	3013	1073	11927 20103 13761 13901 4229 3979	021-170	392 377	-00	1368	27790 41006 26435 32080 11462 13337 4102
	151-160	8 = 5 2 0 0 -	82 82 82	15805 20793 12610 18116 7233 9358 2968	151-160	28.7 28.7 28.0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1117	11749 19823 13448 13600 4228 3979 1134	151-160	319 401 374 340	m w ←	1443	27554 40616 26058 31718 11461 13337 4102
	141-150	¥ 5 5 8 0	3	15709 20679 12540 18073 7233 9358 2967	141-150	238 328	28.	11526 19536 13144 13305 4225 3974 1134	. 81-131	\$ 5 5 8 8	4 + 0	1706	27235 40215 25684 31378 11458 13332 4101
	131-140	200 200 4 - 0	699	15575 20537 12464 17965 7232 9357 2967	131-140	25 28 38 38 4 0 0	1507	11252 19208 12783 13007 4222 3974 1134	131-140	5 23 55 55	0 9 0	2066	26627 39745 25247 30992 11454 13331 4101
	121-130	250 250 128 128 128 128 128 128 128 128 128 128	88	15409 20337 12379 17882 7228 9356 2967	121-130	25 25 4 C O	1688	10939 18785 12400 12825 4218 3972	121-130	52 1 2 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	.00	2363	26348 39122 24779 30507 111446 4101
	111-120	255 271 271 0 0	8	15227 20087 17250 17758 7225 9349 2367	. 81:13	8 25 25 0000	1803	10594 18320 11928 12226 4214 3969 1134	81-11	2 8 8 5 7 8 8 8 8 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9	000	2863	25821 38407 24178 25984 11439 13316 4101
	100-110	7 22 22 23 29 24 2	813	14954 19755 12078 17585 7225 9349 2967	91.90	25 55 55 55 c -	1966	10204 17762 11386 11791 4209 3966 1134	100-110	57.4 87.6 702	22-	280	25156 37517 23464 29376 11434 13315 4101
•	91-100	25 7 25 25 8 26 25 25 8	1471	14730 11922 17388 7218 9339 2967	91-19	52 52 52 5 5 5 6 6 6 6 6 6 6 6 6 6 6 6 6	2778	9654 17205 10626 11266 4197 3963 1133	01-10	85 12 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	25 e	4249	24584 22748 22748 28674 11415 13302 4100
	81-90	55 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	802	14450 19089 11735 17143 7088 9055 2969	8	574 881 735 327 108	2	9396 16534 10292 10761 3677 3691	98:18	979 1454 1054 972	585 4	98	23848 25623 27904 10975 12746 4092
	71-80	207 562 4 51 4 5 1 4 5 1 5 1 5 1 5 1 5 1 5 1 5	98	14045 11416 11416 16756 6906 8764 2956	7-1-80	25.2 25.2 5.5 5.5 5.5 5.5 5.5 5.5 5.5 5.	3803	8624 15653 9557 10176 3550 3383 1132	71-80	1063 1635 1346	è 88 0	828	22869 34129 20973 26972 10458 12147 4088
	61-70	# # # # # # # # # # # # # # # # # # #	3827	13603 17802 10965 16194 6701 8450 2956	61-70	659 1291 949 852 278 278	65	8200 14692 8835 9392 3290 3128 1132	61-70	1341 1536 1538 1691	£ 2.	8117	21806 32494 19800 25586 9991 11578 4088
	51-60	1541 1541 1541 1563 1563 543 966 376	222	12921 16836 16336 15355 6397 8006 2953	81.00	838 1489 1027 1032 310 351	823	2852 2852 2852 2852 2852	8-13	1887 3030 2109 2595	1317	12449	20465 30237 16262 23895 9425 10858 4084
	41-50	1381 2085 1339 2013 1025 1367	9784	11672 15295 9294 13792 5754 7040	2 8	1006 1178 1178 1397 369 463	6497	6706 11912 6859 7508 2718 2501 949	8:	2398 3979 2517 3410	1850	16281	18578 27207 16153 21300 8472 9541 3526
	31-40	2281 2916 1971 3180 1400 1613	14.54	cuba 10481 13210 7385 11779 4778 8653 2033	64.5	1296 2406 1502 1800 467 539 156	9619	111ac	31.40	3557 5324 3503 4980	2352	22322	Otal
	21-30	3848 5018 2840 4175 1718 1963	20018	11.5. Signature 1.5.	, 8 8 8	2061 3706 2043 2177 845 672	11838	NS: St. 4402 7610 7610 1419 1499 625	21:30	5929 8726 4863 6352	2855	82.128 38	126.22 17804 17904 17910 17910 17910 5191 5339 2045
	11.20	2684 3659 2329 3352 1192 1266 466	15148	69 Sun 5276 3344 4424 4424 1611 1657 864	upplied	1500 2548 1435 1555 689 548	9254	2321 3902 2106 2134 1017 2134 313	5.5	4384 6307 3764 4907	645	23782	SSS SSS SSS SSS SSS SSS SSS SSS SSS SS
Ø		1488 1617 815 1072 419 591	6220	ulativ	မ် လ ီ	25 125 125 125 125 125 125 125 125 125 1	4066	11ative 621 539 539 539 538 538	5	2309 2871 1486 1651	352	9	2009 2009 2007 1486 1651 747 870 870 352
Scuba	B.T. — Ough	21 - 28 - 28 - 28 - 28 - 28 - 28 - 28 -	Totals	Cumul.	Surface 11.5	2 2 8 2 8 2 8 2 3 8 2 8 2 8 3 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	Totals	Cumu 21 : 28 22 : 28 23 : 28 24 : 58 24 : 54 25 : 54 25 : 54 26 : 54 27 : 54 27 : 54 28 : 54 2	Total	8 - 38 3 - 38 3 - 38 3 - 38	38 - 18 26 - 18	Totals	Cumu 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2

Figure A1

All Dives




Table A2Distribution of Dives by Depth and Bottom Time

Beryo

1 8 2 4 5 E

e Beyo

• Beryo No-D LI

84384 2

Navy Dives

Totale 7 12267 16141 16141 14441 5845 8877 2136		Totale h 13422 113422 113422 113422 113424 8854 8854 8854 8854 8854 8854 8854 8	\$563 \$	1 Totale N Totale N 222215 29643 20726 21297 8669 8669 8669 3218 3218 3218	## T T T T T T T T T T T T T T T T T T
381.360	12267 16141 10626 14441 5845 6977 2336	25.138	9 9948 13422 10200 6856 1704 1704	25. 86. 80.0000	22215 22215 20726 21277 8669 8681 3218
141.350	12264 16138 10526 14441 5845 6977 2336	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1 13421 13421 10195 6856 6856 1704	. 1	3 22209 29559 20721 220721 8669 8681 3218
231-340	12264 16137 10525 14441 5845 6977 2336	231.240	1 9945 13420 10195 6856 8856 1704	23.540	13 22209 29557 20720 20720 8669 8681 3218
25.130	12262 16137 10518 14440 5845 6975 2336	. 200	1 9945 13420 10194 6856 2824 1704	25. 4 9 9 9 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	19 22207 29557 20712 21296 8669 8679 3218
311.220	12259 16128 10512 14440 5845 6975 2336	25.	14 9944 13420 10134 10134 1704 1704	21.220	26 22203 22203 220706 21296 8659 8679 3218
200.73.00	12252 16125 10512 14438 5845 6975 2336	301.210 84 8 0 0 0 0 0	129 9930 13420 10194 6656 2654 1704	304.310	148 22162 20706 21294 8669 8679 3218
29. 300	12250 16116 10506 14436 5843 6975 2336	281-300 33 33 55 75 90 0 0 0 0	9900 13372 10143 6856 2824 1704 8872	291.300 8.80 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	245 22150 220649 21294 8667 8679 3218
13 13 13 13 13 13 13 13 13 13 13 13 13 1	12246 16106 10482 14438 5843 6975 2336	281-290 20 20 73 73 65 65 0 0 0 0	169 9867 13273 10068 6856 2824 1704	281-290 42 80 6 6 0 0	206 22113 22650 21254 8667 8679 3218
271-280 8 8 27 20 0 0	12233 16093 10469 14434 5843 6975	271-280 37 63 51 51	9436 13200 10003 9854 2624 1704	271-280 45 71 78 6 0	200 22071 29299 20472 21288 8657 8679 3218
261-270 25	12225 16091 10442 14428 5843 6975 2336	261.270 38 76 80 0 0 0	195 9952 6854 1704 862	28 8 8 5 0 0 0 0	250 22026 29228 20394 21382 8667 8679 3218
251-260	122.10 16072 10423 14426 5843 6975	251-260 44 89 76 0 0	209 9762 13061 9872 6854 1704	251-260 55 1113 97 0	265 21972 220295 21280 8667 8667 3218
241-250 28 38 6 0 0 0 0	12199 16048 10402 14426 5843 6975 2336	24-256 38 8 8 73 8 8 9 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	208 97.18 12972 9796 6854 2824 1704	241-250 38 106 6 0 0	257 21917 22020 21260 8657 8657 8679 3218
231-240 255 36 36 0 0	12194 16036 10376 14420 5843 6975 2336	231-240 73 117 102 0 0	301 9685 12876 9717 6854 2824 1704	231-240 104 166 138 0	410 21879 20093 21274 8667 8679 3218
22. 24. 25. 20. 00. 00. 00. 00. 00. 00. 00. 00. 00	12169 15993 10340 14415 5843 6975 2336	221-230 88 105 105 0	253 9606 12759 9615 6452 2824 1703	221:230 88 8 8 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	303 21775 28752 21267 8667 8678 3218
21-220 22 22 20 19 0 0 0 87	12155 15969 10330 14413 5843 6975 2336	93 93 126 127 127 0 0	348 8540 12654 8533 6852 2824 1703	211-220 115 115 126 10 0	426 21696 226623 19653 21265 8667 8678 3218
201-210 20 20 20 20 20 20 20 20 20 20 20 20 20 2	12133 15940 10311 14405 5843 6975 2336	201-210 102 140 82 3	337 9447 12528 9408 6850 2824 1703	201-210 138 202 202 132 6 6	465 21580 28468 19717 21255 8667 8678 3218
191-200 42 43 46 0 0 0 0	12100 15878 10281 14402 5843 6975	191-200 116 175 185 148 0 0	624 9345 12388 9314 6847 2824 1703	158 218 213 0 0	783 21445 28266 19595 21249 8667 8667 3218
181-190 51 28 28 0 0 0 0	12058 15835 10253 14356 5843 6975 2336	181-190 132 239 157 196 0 0	724 9229 12213 9129 6699 2624 1700 862	178 290 118 234 234 0 0	874 21287 28048 19382 21055 8667 8678 3218
171-180 121 121 121 121 121 121 121 121 121 12	12012 15784 10228 14328 5843 6975 2336	171-180 169 247 222 173 173 0	9097 11974 8972 6503 2824 1703	171-180 239 368 256 256 9 9	21109 27758 27758 19200 20831 8667 8678
52 52 50 50 69 60 0	11942 15663 10194 14265 5637 6974 2336	158 158 234 200 133 0	728 8926 11727 8750 6330 2621 1703 862	161-170 210 333 253 182 0	978 20870 27390 18944 20595 8658 8677 3218
151-160 85 91 56 40 0 0 0 1	11890 15564 10144 14216 5837 6974 2336	151-160 190 223 204 137 2 2	8770 11493 8547 6197 2821 1703 862	275 314 262 177 177	20660 27057 18691 20413 8658 8677
105 105 126 68 68 63 10 10 10 10 10 10 10 10 10 10 10 10 10	11806 15473 10066 14176 5837 6974 2335	141-150 228 2861 224 224 0	8580 11270 8343 6080 2819 1701	141-150 333 387 292 194 194	20385 26743 18429 20236 8656 8675 3217
131-140 138 163 71 65 65 65 4 4	11700 15347 10018 14113 5836 8973 2335	131-140 244 329 250 250 156 0	8352 11009 8119 5929 2817 1701	390 422 321 223 7 7	1424 20052 26356 18137 20042 8653 8674 3217
121-130 160 196 98 87 7 7 0	11564 15184 9947 14048 5832 6972 2335	121-130 256 333 296 165 4 4	1057 8108 10680 7869 5771 2814 1701 882	-	19672 25864 17816 19819 8646 8673 3217
233 238 258 117 117 134 0	11404 14996 9849 13968 5830 6965 2335	280 280 370 351 193 4	7852 (0347 7573 5606 2810 1698	-	1955 19256 25345 17422 19574 8640 8663 3217
100-110 134 134 0 0 0 727	11171 14740 9732 13834 5830 6965 2335	250 250 371 233 4	7559 9977 7222 5413 2806 1696 882	446 619 505 505 175 10	1958 18730 24717 16954 19247 8636 8636
219 274 140 172 77 77 8	10975 14493 9598 13692 5824 6963 2335	91-100 310 411 411 411 525 236 236 236	7309 9605 6851 5180 2802 1696 881	91-100 529 685 462 462 308 184 184	2604 18284 24096 16449 18872 8626 8626 8659 3216
81-90 303 416 273 273 152 90 3	10756 14219 9458 13514 5752 6886 2327	81-90 347 513 461 271 223 107	1963 6999 6194 6529 4930 2566 1569	61-90 690 929 975 183 375	3410 17755 23413 15967 18444 8318 8475 3208
71-80 337 4114 402 129 0	10453 13603 9238 13241 5600 6806	71-80 442 528 528 444 346 170 0	2006 6612 8661 6068 4659 2343 1482 860	71-80 779 942 860 748 331 199	1798 22484 15306 17900 7943 8288 3204
61-70 480 613 470 612 235 235 231 244		61-70 637 540 540 160 160 1	2197 6170 6170 8153 5624 4313 2167 1412 860	61-70 933 1250 1010 955 395 294 294	16286 21542 114506 17152 7612 8089 3204
51-60 748 11119 824 1236 521 734 264 5445	9636 12778 8412 12227 5210 6446	51-60 596 796 796 410 196 196	2818 5717 5718 5084 3970 2007 1349 1349	51-80 1344 1915 1448 707 841	15353 20292 13496 16197 7217 7217 7200
41-50 924 1464 1065 1560 813 1074 446	CC .	41-50 732 872 880 880 486 214 125	Ö	41-50 1656 2336 1735 1027 1199 560	10558 14009 18377 12048 14551 6510 6954 2837
31-40 1602 2160 1572 2465 1151 1427 486	Scub; 7864 10193 8533 8431 4636 1611	31-40 919 1120 677 648 262 262	Surfa 4389 5848 3074 1607 1117 666	31.40 2521 3280 2449 3113 1413 1591	14977 Total 12353 16041 10313 12505 5483 5755
21-30 2825 3900 2334 1406 1406 1597 569	=	0	24.78 24.78	21-30 4402 6012 3692 4278 1997 1937	23164 JMS: 9832 12761 7864 9392 4070 4164
11.20 2275 2794 1902 2790 960 1086 358	>	Suppli	Ye Size 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2		8
10a 1242 1242 1242 1244 1244 1244 1244 124	_	Surface Sur part 2 200 100 100 100 100 100 100 100 100 1	3		100 100 100 100 100 100 100 100 100 100
Scuba	2 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	LE 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	2 2 8 2 8 2 8 2 8 2 8 2 8 2 8 2 8 2 8 2	5. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.	Cum (2 % % % % % % % % % % % % % % % % % %

Figure A2 Navy Dives




Table A3Distribution of Dives by Depth and Bottom Time

	U	D	
	d	٥	
	>	>	
•	Ξ	-	
ĺ)	
	2	-	
	ā	3	
_	C	S	
:	=	=	
	=	-	
	7	>	
2		=	
(_)	
	_		

!	No-DLi	•	# Beyo	0.270	n	e Beryc No-D Li	0.220 2	
	Totale 1213 2813 1276 2153 732 1608	8	Total	2718 8022 5109 7804 1335 2214 198	27461	Totale	3932 10838 6385 8957 2067 3422 503	
	351.360	0 1213 2813 1276 2153 732 732	86.18	• • • • • • • •	0 2719 8022 5109 7804 1335 2214	361.360		3932 10835 5385 2067 3922 503
	341.350	0 1213 2813 1276 2153 732 1608	¥ 550	000000	0 2719 8022 5109 7804 1335 2214	341-350		3932 10835 9957 2067 503
	331.340	0 1213 2813 1276 2153 732 1608			0 2719 8022 5109 7804 1335 2214			3902 10035 6385 9957 2067 3822 503
	000000	0 1213 2813 1276 2153 732 1608	0	000-000	1 8022 5109 5109 1335 2214 198	•	000-000 -	3932 10835 6385 9957 2067 503
	11.320	1213 2813 1276 2153 732		N F1 0 0 0 0 0	\$ 2719 8022 5109 7303 1335 2214 198		nn00000 w	3932 10835 6.385 9956 2067 503
	301-310	1212 2813 1276 2153 732		m + 1 0000	16 2717 8019 5109 7803 1335 2214 198		w+-\0000 #	3929 10832 6385 9956 2067 3822 503
	291,300	1212 2813 1276 2153 732 732		01100+0	10 2712 8015 5102 7803 1335 2214 198		00-00-0 9	3924 10628 6.378 9956 2067 503
	281.290	0 1212 2813 1276 2153 732 1608		NO 0 - 000	8 2712 8013 5095 7803 1335 2213 198		NO 0 - 0 0 0 0	3924 10826 6371 9956 2067 3821 503
	271.260	1212 2813 1276 2153 732 1608		#WN00-0	2710 8013 5092 7802 1335 2213		nun00-0 =	3922 10026 6368 9955 2067 3821 503
	261-270	0 1212 2813 1276 2153 732 1508	261.270	* 10 m	13 2707 8008 5090 7802 1335 2212 198		4 N m - 000 5	3919 10821 6366 9965 2067 3820 503
	13 13 0	16 1212 2813 1278 2153 732 1608		- 80000	20 2703 8003 5067 7801 1335 2212 198	•	4 2 5 0 0 0 0 %	3915 10816 6363 9954 2067 3820 503
	241-250	1209 2800 1276 2153 732		: :: 0000	22 2702 2702 7997 5074 7801 1335 2212		5 0000 X	3911 10797 6350 9954 2067 3820 503
	231-240	16 1209 2797 1276 2153 732 1608		m r o = 0 0 0	20 2669 7994 7994 1335 2212 198		2 000 S	3898 10731 6344 9954 2067 3820 503
	221-230	11 1200 2786 1276 2153 732		να. <mark>‡</mark> 0000	22 2686 7967 7967 7800 1335 1335 198	0	~2₹0000 B	3886 10775 6335 9953 2067 3820 503
	211-220	1195 2782 2782 1276 2153 732		• 2 • n o o o	31 2684 7961 5045 7800 1335 2212 1396		# C 0 0 0 E	3879 10763 6321 2953 2067 3820 503
	201-210	1196 2782 1276 2153 732		0000 25 2 0	51 2675 7971 7797 1335 2212		8 000 825	3870 10753 6312 2067 3820 503
	191.200	31 2768 1276 2153 732 1608		2832	192 2666 7957 7795 1335 2212 196		28 × 2000 8	3360 10725 6286 9948 2067 3820 503
	2 2 9 9 7 7 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	24 1193 2759 1269 2139 732 1608	11.190	# 2 2 2 E	2654 7928 7708 1335 2212	061-190	8 % <u>2 %</u> 000 <u>8</u>	3847 10687 6215 9847 2067 3820 500
	171-180 15 15 15 15 15 15 15 15 15 15 15 15 15	37 1191 2750 1262 2133 732 1608	171-180	225	2626 7099 4844 7528 1335 2212	171-190	8 4 ± 5 0 0 0 5	3817 10649 6106 9661 2067 3820 503
	161-170 13 14 9 0	40 1183 2735 1253 2128 732 1608	021-170	8152-00	343 2596 7865 4712 7357 1335 2212	161-170	124 174 174 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	3781 10600 5965 9485 2067 3820 503
	151-160 8 23 12 4 4 0 0 0	1179 2722 1239 2119 732 1508	151-160	83872-00	355 2578 7821 4602 7189 1334 2212	151-160	88 = = = 0 00	3757 10543 5841 9308 2086 3820 503
	141-150 133 73 23 0	2699 1227 1227 2115 732 1608	141-150	\$ \$ \$ \$ - 0 0	2546 7758 4503 7032 1333 2209	141-150	2 8 1 5 0 0 C C C C C C C C C C C C C C C C C	3719 10457 5730 9147 2065 3817 503
	131-140 23 34 33 33 0 0	1157 2686 1220 2092 732 1608	131-140	88222-00	2501 7691 4367 6865 1332 2209 196	131-140	257 127 127 127 127 127 127 127 127 127 12	3658 10377 5587 8957 3817 503
	21.130	153 2652 1208 2059 732 1608	121-130	\$ 5.50 0 0 0 0 0 0	2433 7596 4239 6641 1331 2207	121-130	8 8 8 8 - 0 0 ET	3567 10250 5445 8700 2063 3815 503
	11-120	185 2592 1176 2016 731 1608	111-120	8 5 5 5 0	2347 7468 4069 6406 1331 2207	111-120	2250 234 234 234 234 234 234 234 234 234 234	3462 10060 5245 8424 3415 503
	100-110 23 58 19 19 0	156 2527 1127 1979 731 1608	100-110	7.6 1.75 7.75 0.0 0.0	740 2252 7283 3884 6167 1330 2206 196	100-110	120 238 201 319 7 7 11 1 1 0 0	3333 9610 5011 2061 3614 503
	81.58 82.88 82.00 82.00 83.00 80 80 80 80 80 80 80 80 80 80 80 80 8	450 1058 2469 1108 1931 731 1600	91-100	4 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	2155 7103 3702 5896 1323 2203 198	91-100	189 309 251 336 125 367	3213 9572 4810 7827 2054 3803 503
	81-90 85 85 85 85 102 102 0	2415 1003 1003 1869 6390 1397	91-90	25.7.7.2.2.7.7.2.0.0 0.0000000000000000000	2011 6848 3496 5622 1239 2039	9.	240 515 361 415 129 412 0	3024 9263 4559 7491 1929 3436 503
	71-80 88 212 88 133 40 0	967 2228 973 1767 662 662	71-80	52 22 22 22 22 22 22 23 25 30 0	1827 6489 3225 5309 1138 1836	71-80	236 360 360 173 173 359 0 0	2784 6748 4198 7076 1900 3024 503
	61-70 108 279 101 187 58 200 0	803 891 2047 887 1634 622 1012	61.70	8 2 8 8 5 5 °	2951 1657 1965 1965 1965 1965 1965 1965 1965 1965	61-70	286 286 287 288 288 288 288	2548 8107 3838 5508 1677 2665 503
	51-60 28-68 28-61	1062 783 1768 786 1447 564 812	81-60	228 6480 397 609 123 241	2551 1467 1467 2551 2551 1647 1647	81-60	318 946 557 203 433 157	2250 7167 3337 5818 1518 2253 503
	41-50 320 134 271 108 197 62	CO CO	•	261 995 476 901 174 45	5	8:14	408 1315 610 1172 282 533 107	1932 6241 2780 5011 1315 1820 346
	31-40 155 330 142 345 167 167 45	Scubing 1280	31-40	330 1221 638 1136 368 28	Surfa 274 274 2861 2861 2861 2861 2861 2861	31.40	485 1551 780 1481 302 535 73	Total 1524 4926 2170 3839 1033 1287 239
	21-30 192 395 175 345 128 128 151		ied 2:30	341 1425 577 1080 232 312	2522 2523 2623 2623 2633 2633 2633 2633	21-30	533 1820 752 1425 360 463 66	1038 1038 3075 1390 2358 731 752 166
	11-20 148 354 147 224 115 91	Ve Su 188 188 188 188 188 188 188	Jddn	21, 25 88 2(9) 21, 25 88 2(9) 31, 35 40 207 389 31, 35 40 27 389 41, 45 58 171 46, 50 26 158 51, 55 4 22	8	11:20	367 1245 527 772 286 249 86	Ve Su 506 1555 608 933 371 283 100
þa	0-10 103 103 103 103 103 103 103 103 103 1	3	9 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	207 207 4	3	_	139 310 111 161 161 14 85 85	138 339 339 339 339 339 339 339 339 339 3
Scuba	######################################	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	Surf J. J. J	2888282 2888383 2883383	2	Total	28 - 28 21 - 28 22 - 38 23 - 36 24 - 45 25 - 55 27 - 55 27 - 55 28 - 55 27 - 55 28 - 5	2 8 8 2 8 2 8 2 8 2 8 8 2 8 2 8 2 8 2 8

Figure A3

Civilian Dives




Table A4

Distribution of Dives by Depth and Bottom Time

/۸	
les	
5	
<u>Ф</u>	
듣	
Ma	





Figure A4 Marine Dives



36 - 40

26 - 30 - 31 - 35 -

21 - 25 🖃

51

46 - 50 --

- 45

Table A5


Distribution of Dives by Depth and Bottom Time

Reserves Dives

,	
•	
,	
)	
•	

Scuba

# Bey No-D	••••	۰	
Totale	1392 1501 747 1260 459 505	\$648	
351-360		۰	1392 1501 747 1260 459 505 181
341-350		۰	1392 1501 747 1260 459 505 181
331-340		۰	1392 1501 747 1280 459 505 181
321-330		۰	1392 1501 747 1260 459 505 181
311-320		0	1392 1501 747 1260 459 506
301-310		0	1392 1501 747 1260 459 181
291-300		۰	1392 1501 747 747 459 459 161
281-290		•	1392 1501 747 747 1260 459 161
271-280		0	1392 1501 747 1260 459 506
261-270		0	1392 1501 747 1260 459 505 181
251-260	+000000	•	1392 1501 747 1260 459 505
241-250		0	1388 1501 747 1260 459 505
231-240	00-000	r	1388 1501 747 1260 459 506
221-230	000000	0	1388 1499 746 1260 459 505
211-220	0000000	0	1386 1499 746 1260 459 505
201-210		0	1388 1499 746 1260 459 505
191-200	*****	•	1388 1499 746 1260 459 505
181-190	00-000	-	1384 1499 746 1260 459 506
171-180	noo-ooo	e	1384 1499 745 1260 459 505 181
161-170		•	1382 1499 745 1259 458 505
151-160	*000000	•	1381 1499 745 1256 459 505
141-150	₽n-4000	6	1377 1499 745 1256 459 505
131-140	8 700000	2	1384 1496 744 1254 459 459 181
121-130	mm000		1358 1493 744 1249 459 459 181
111-120	88-NO00	10	1355 1490 743 1248 459 505 181
100-110	R 2 4 7 0 0 0	28	1349 1481 742 1246 459 505 181
91-100	21001-0	8	1344 1468 739 1239 458 505 181
91-90	\$5 8 5 8 0 O	2	1330 1461 737 1234 452 504 181
71-80	£802500	3	1312 1445 728 1222 443 504
61-70	88 82 58 → £5 o	142	1295 1419 719 1201 431 485 181
51-60	848858	278	1256 1381 706 1166 427 482
41.50	8 5 5 7 2 8 2 7 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	583	1200 1337 654 1103 412 452 163
31-40	257 248 150 100 128 44	1167	326 338 388 388 388 388 388 388 388 388 38
21:30	472 486 223 416 126 168	1952	E 25 25 25 25 25 25 25 25 25 25 25 25 25
11-20	272 325 155 259 259 77	1216	76 St. 252 252 252 252 252 252 252 252 252 25
0-10	8 ≩ 3 2 2 2 5 5	\$	ulative
₩.T.	2 2 8 2 8 2 8 2 2 2 8 2 8 2 4 2 8 5 6 6 6 6 8 8 8	Totals	2







Figure A5

Reserves Dives



36 - 40 26 - 30 - 31 - 35 -- 55 51 46 - 50 --21 - 25 -41 - 45 --

APPENDIX B

Part I. Summary of Cases

Code

Purpose of Dive

SW = SPECWAR

SH = Ship Husbandry

TN = Training

UC = Underwater Construction

SR = Search

SS = Security Swim

EO = EOD

SV = Salvage

RQ = Requal

IS = Inspection

Rig

SO = Open circuit SCUBA

SS = Surface-supplied

NMRI Diagnosis

Class

1 = Definitely Decompression Illness

2 = Definitely not Decompression Illness

3 = Uncertain if Decompression Illness or not

SubClass

A = Decompression Sickness

B = Arterial Gas Embolism

C = Decompression Sickness or Arterial Gas Embolism

N = Not applicable

Naval Safety Center Diagnosis (SAFE Dx)

1 = Type I Decompression Sickness

2 = Type II Decompression Sickness

3 = Arterial Gas Embolism

Service

N = Navy

M = Marine

C = Civilian

Part II. Case Descriptions

Narratives exactly as received from NSC; "(sic)" used to denote obvious and/or distracting errors Cases with bottom times longer than the USN93 limit are indicated with an asterisk.

Part I. Summary of Cases

		vi.ii								Diagnosis		
Case		ay	Month	Year	Purpose	Rig	Depth	Bottom Time	Class	Subclass	SAFE Dx	Service
1	133. SU			90	SW	SO	50	52	3	N	2	. N
2 3*	oran er Sauts	3 2	2	90	SW	SO	50	52	3	N	2	N
4*	or broad filter	_		90	SW	SO	30	397	3	N	2	N
4		16 21	3	90 90	SW TN	SO	30	397			2	N
6	Ludela, Subs	18	4	90	TN	SO	52 45	13	1	. A	2	N
7			6	90	UC	SS	52	46 46		A	1	M
8	filling States a	7	8	90	SH	SO	40	74	1	A A	2	N C
9		29	, ,	90	SW	so	29	135		A	2	N
10		23	10	90	SS	SO	36	30	3	N	2	N
11*		6	11	90	SH	SS	40	189	ī	A	2	N
12		5	12	90	SH	SS	26	50	3	N	2	N
13		15		. 91	SR	SS	55	29	2	N	2	N
14		27	1	91	SH	SS	27	17	1	C	2	N
15	9 WY W H	1	2	91	SH	SS	30	30	1	С	2	N
16		11	2	91	SH	SS	37	134	1	Α	1	N
17		5	4	91	SH	SO	35	76	1	С	2	Ň
18		16	2	93	SH	SS	31	87	2	N	2	N
19		23	. 5 3	93	SH	SS	41	99	2	N	2	N
20		29	3	93	SH	SS	42	97	3	N	2	N
21*		21	7	93	SW	SO	33	278	1	ΑΑ	1	N .
22	er Leverine granning	9	8	93	SH	SS	25	140	3	N	2	N
23	a kadi Dik	16	. 3	94	RC	SO	30	45	1	Α	2	N
24	vili ya sana asanini	21	4	94	SH	SS	22	91	3	N	2	N
25	teliki 1894	27	5	94	SH	SS	38	91	1	C	2	C
26	21 1 2 10 Andrews	12	. 7	94	SW	SO	28	265	2	N	2	N
27	a an i iiria.	. 8 2	8	94	SH	SO	45	15	3	N	. 2	N
28 29	aga provent ing province	8	8	94	SH	SS	36	113	<u>1</u>	A	2	N
30	YFRE S SIGN	17	8	94	SR	SO	27	73	3	N	2	N.
31	www.warana	29 14	2	90 90	TN SW	SO	22	29	1	C	3	N
32	168 (197	24	4	90	TN	SO SO	40	121	. 1	· . C	3	N
33	arest take	21	100003	90	SW	SO	24 36	40 50	2	A N	3	N
34	visi i Winne	21	5	90	SW	SO	36	50	3		3	N
		19	7	90	SR	SS	26	61	3	N C	3 3	N
36*	PPONE ARREST	21	8	90	SH	SS	37	176	1	A	3	N N
37			10	90	SH	SS	25	44	1	iii.c	3	N N
38		12	11	90	SV	SS	33	134	3	N	3	N
39*		9		91	SH	SS	34	191	3	N	3	N
40		4	4	91	SR	SO	40	3	2	N	3	N
41		30	5	91	SV	SO	35	52	. 1	C	3	N
42		30	5	91	SV	SO	35	52	3	N	3	N
43 : . :				91	EO	SO	50	40	. 1	A	2	N
44	1197 84421.	16	7	91	SH	SS	26	25	1	C	3	N
			8	91	TN	SS	29	22	3	N	3	N
46	in the second	5	9	91	SW	SO	29	74	1	C	3	N
				91	SH	SS	35	115	3	N	3	N
48 49	165 703	6	11	91 93	EO	SO	45	23	1	В	3	N
50	. R'4 + 11(+ 2)				IS	SO	47	57	1	C	3	С
51		25 16	2	93 93	SS RC	SO SO	40	20	1	В	3	N
52		21	3	93	SV	SS	42 52	20	. I	В	3	N
53		27	7	93	SR	SO	22	58 25		C	3	N
54		31	8	93	SH	SS	28	37	2.	N	3 .	N
55		13	10	93	SR	SS	35	43	1	B	3	N
56		18	11	93	IS	SO	23	43	1	A	3	N
57		28	rei i	94	EO	so	37	20	3	A N	3 3	N
58		23	2	94	EO	SO	40	20	1	В	3	N
59		25	3	92	SW	so	25	40	1	A A	3	N N
60		30	8	92	SH	SS	21	63	Î	A	2	N N
								95	•	73	4	IN

Part II. Case Descriptions

Case 1

This 27 year old male spec war diver made a no "D" dive using SCUBA. The dive was on air to 50 fsw for :52. Upon surfacing the diver experienced extreme fatigue following submarine lock out. The diver was examined and no other deficit was found. He was compressed to 60 fsw and had complete relief :55 at depth. The diver was treated on a TT 6 with one extrnsion (sic) at 60 fsw. No meds were used, final diagnosis was DCS II, treated by DMO, 2 days loss of work and 28 days restricted from diving.

NMRI Diagnosis: Uncertain if decompression illness or not

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 2

This 25 year old male Spec War diver made a no "D" dive using SCUBA. The dive was on air to 50 fsw for :52. Upon surfacing the diver experienced extreme fatigue following submarine lock out. The diver was examined and no other deficit was found. He was compressed to 60 fsw and had complete relief :55 at depth. The diver was treated on a TT 6 with one extension at 60 fsw. No meds were used, final diagnosis was DCS II, treated by DMO, 2 days loss of work and 28 days restricted from diving.

NMRI Diagnosis: Uncertain if decompression illness or not

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 3*

This 40 year old male Spec War diver made an SDV dive using open circuit SCUBA. The dive was to 30 fsw for 6 hours and 37 mins. The dive was reported as a no "D" dive. According to standard air tables the dive was 40 for 480 exceptional exposure because the max time of the repetative (sic) group designator table was exceeded. After a restless night the diver reported to work in the morning. Physical discomfort was reported to the diving supervisor. A neurological exam was normal but neuropsych screening was borderline. The diver was compressed to 60 fsw and treated on a treatment table 6. No meds were used. No days loss of work, 4 days restricted from diving, final diag. DCS Type Type (sic) II by DMO. The other diver on this dive had DCS II and treated (sic).

NMRI Diagnosis: Uncertain if decompression illness or not

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 4*

This 41 year old male Spec War diver make (*sic*) an SDV dive using open circuit SCUBA. The dive was to 30 fsw for 6 hrs and 37 mins. The dive was reported as a no "D" dive. According to standard air tables the dive was 40 for 480 exceptional exposure because the max time of the repetitive group designator table was exceeded. The following morning the diver reported a restless evening and a physical discomfort to the Div Sup. A neuro revealed decreased light touch and pin prick sensation over right lateral thigh. The diver was compressed to 60 fsw and treated on a TT 6. No meds were used, no loss days from work, 4 days restricted from diving. Final diag. DCS II by DMO. The other diver on this dive had DCS II and treated (*sic*).

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

This 29 year old Navy male 1st class diving student made a training dive using Mark XII SSDS air. The dive was to 52 fsw for :13 min bottom time no "D". Forty five minutes after surfacing he reported pain in the left shoulder and elbow. A neuro also revealed nystagmus in the right eye. The diver was compressed to 60 fsw and treated on a TT 6 with 2 extensions at 60 fsw. Complete relief was noted at 8 mins. into 2nd O2 period at 60 fsw. Medications used: 1000 cc of normal saline. Two days loss of work and 14 days restricted from diving. Final diagnosis DCS Type 2 by MDV.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 6

This 20 year old male Marine Corps SCUBA student made an open water SCUBA training dive. The dive was to 45 fsw for :46 min no "D". Nine hours after surfacing the diver complained to the supervisor of pain in the left knee. A neuro was conducted and neurological deficit was found. The diver was compressed to 60 fsw and treated on a TT 6. Complete relief occurred during the 4th O2 period at 60 fsw. No drugs were used, no days loss from work and 28 days restricted from diving. Final diagnosis was DCS Type one by DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

Naval Safety Center Diagnosis: Type I Decompression Sickness

Case 7

This 21 year old male 2nd class diver made an underwater coinst. ops dive in open water using MK XII, air, swimmers configuration. The depth was to 52 fsw for 46 min no "D". Approx. five minutes after surfacing the diver noted a dull pain behind right kneecap. A neuro was completed by a DMT with no other findings. The same pain waxed and waned over the next 50 minutes. The diver was compressed to 60 fsw, complete relief was obtained upon reaching treatment depth. Treated on a treatment table 5, no days loss of work, 5 days restricted from diving, no meds used, final diagnosis DCS I by DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

Naval Safety Center Diagnosis: Type I Decompression Sickness

Case 8

This 26 year old male ship yard diver made a ships husbandry dive using open circuit SCUBA. The dive was to 40 fsw for 19 minutes no "D". The previous day this diver made a no "D" dive to 40 fsw for 74 minutes bottom time. During ascent of this dive he noticed pain in his right hip. This pain was not reported. The next dive he made the 40 fsw for :19 dive. He noticed the pain improved on the bottom of this dive. On ascent this pain increased. Upon surfacing he was taken to the DMO and compressed to 60 fsw and had complete relief upon reaching treatment depth. Due to the location of the pain he was treated on a TT 6 with one extension at 60 fsw. No meds were used, one day loss of work, 14 days restricted from diving, final diagnosis was DCS II by DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

This 24 year old male SEAL made an SDV training dive with SDV air system and full face mask. The gas media was air. The dive was to 29 fsw for 2:15 no "D". :45 after surfacing the diver reported numbness to left nostril and left quadriceps. He was compressed to 60 fsw and had complete relif upon reaching treatment depth. No meds were used, no days loss of work, 14 days restricted from diving. Final diagnosis was Type 2 DCS by DMO. Suspect circulation impaired by tightness of equipment.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 10

This 24 years old male SCUBA diver made a security swim using open circuit SCUBA. The dive was to 36 fsw or (sic):30 BT. Approx 3 hours after the dive this diver became dizzy, felt nauseated and had a painful left ear. A neuro was performed by a DMO and DCS Type II was diagnosed. The diver was transported to a recompression chamber and compressed to 60 fsw on O2. Complete relief was obtained during the 1st O2 period. He was treated on a TT 6, Meclizine was given, no days loss of work, 30 days restricted from diving.

NMRI Diagnosis: Uncertain if decompression illness of not

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 11*

This 23 year old 2nd class diver made a ships husbandry using MK 21 air. The dive was to 40 fsw for 180 minutes no "D". Approx one hour after surfacing the diver noticed pain and a slight numbness in his left forearm. The diver did not report these symptoms until five hours later. He was compressed to 60 fsw on O2 and had 1st relief on descent at 30 fsw. Complete relief was obtained 33 minutes into the treatment. A TT 6 was completed, no meds were used, no days loss of work and 14 days restricted from diving. Final diagnosis DCS Type II by DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 12

This 19 year old 2nd class diver made a ships husbandry dive using MK 21 air. The dive was to 26 fsw for :50. No "D". Upon surfacing this diver reported that his foot was numb. A neuro was completed with no significant findings. He was transported to a chamber and treated on a TT 6. First relief was noted :13 into the treatment and complete relief at :29. No meds were used, no days loss of work and 21 days restricted from diving. Final diagnosis was DCS II by DMO. Phoncon to his master diver, who was TAD at the time of the dive and the treatment, reported that this diver was very cold and tired and wore an ill fitting wet suit. He feels that the DCS II diagnosis was highly unlikely due to the dive profile.

NMRI Diagnosis: Uncertain if decompression illness or not

This 33 year old Navy Saturation Diver made a search dive using MK 21. The dive was to 55 fsw for :29 BT no "D". Upon surfacing the diver noticed numbness and tingling in the top of his left foot. The diver also reported that his boot was too tight. MDV and DMT performed thorough neurological examination. No other findings were noted. DCS could not be ruled out so the diver was compressed to 60 fsw on O2. Partial relief of symptoms were noted after :03 at depth and complete relief was obtained after :14. No meds were used, no days loss of work and 14 days restricted from diving. Final diagnosis DCS Type II by MDV and DMT.

NMRI Diagnosis: Definitely not decompression illness

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 14

This 32 year old Navy HeO2 diving officer made a ships husbandry dive using MK-1 air. The dive was to 27 fsw for :17 BT. :67 after surfacing the diver complained of left arm tremors, numbness, and weakness. Deamethasone (sic) 10 MGM was given during medivac to the recompression chamber. He was compressed to 60 fsw on O2. First relief was obtained upon reaching depth and complete relief during the first O2 breathing period. He was treated on a TT six. Three days loss of work. 28 days reftricted (sic) from diving. Final diagnosis DCS II.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness or Arterial Gas Embolism)

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 15

This 28 year old Navy 1st class diver made a ships husbandry dive using MK-1 air. The dive was to 30 fsw for :30 no "D". Approx 15 min after surfacing the diver felt slight disorientation but did not report this until 40 mins later. A neuro exam revealed vestibular imbalance and lower extremity weakness. He was placed on O2 and transported to recompression chamber. Compression was to 60 fsw on O2. Complete relief was obtained 22 minutes into the treatment. A TT 6 was completed. No meds were used, one day loss of work and 7 days restricted from diving. Final diagnosis was DCS Type II by DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness or Arterial Gas Embolism)

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 16

This 20 year old male 2nd class diver made a ships husbandry dive using ESDS AGA. The dive was to 37 fsw for 2:14 no "D". Approx. four hours after the dive the dive (sic) the diver complained of pain in the right thigh. No type two symptoms were noted. The diver was compressed to 60 fsw and had first relief eight minutes into the treatment. Complete relief was noted twenty one minutes later. A TT 6 was completed. No meds were used, no days loss of work, 7 days restricted from diving. Final diagnosis, DCS 1 by MDV.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

This 30 year old 1st class diver made a ships husbandry dive using open circuit SCUBA. The dive was to 35 fsw for 1:16. Thirty minutes after surfacing the diver complained that his left hand was not feeling as it should. A neuro exam showed numbness of the left thumb, index, and middle finger. He was compressed to 60 fsw on O2. First relief was noted upon reaching treatment depth and complete relief:09 into the treatment. A TT 6 was completed. No meds were used, no days loss of work and 14 days restricted from diving. Final diagnosis was DCS II by DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness or Arterial Gas Embolism)

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 18

This 22 year old male 2nd class diver made a ships's (*sic*) husbandry dive using MK 21. The dive was to 31 fsw for 8M (*sic*) mins. Three weeks after dive this diver reported severe headaches, dizziness, and memory loss. Neuro exam conducted by DMO and MDV with no neuro deficit noted. Referred member to Naval Hospital for full neurological testing and decision made to treat member on htperbaric (*sic*) oxygen. Recompressed to 60 fsw with no relief noted during treatment. A TT6 was completed. No meds were used. 17 days loss of work and 17 days restricted from diving.

NMRI Diagnosis: Definitely not decompression illness

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 19

This 26 year old male first class diver made a MK-21 dive to 41 fsw for 99 mins while inspecting patches for leaks during salvage operations on a ship. 4 hrs 50 mins after surfacing diver complained of numbness to left great toe. Neuro exam conducted with no other symptoms noted. Diver recompressed to 60 fsw and treated on a TT-6 for DCS type II. Diver had complete relief 4 hrs 37 mins into the treatment. No meds were used. No days loss of work and 14 days restricted from diving.

NMRI Diagnosis: Definitely not decompression illness

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 20

This 21 year old male second class diver made a ship's (*sic*) husbandry dive using ESDS to a maximum depth of 42 fsw for 1 hr 37 mins. Approximately 10 hrs after surfacing diver reported radiating pain in his left leg and "something not being quite right". Neuro exam revealed numbness on the left leg from the knee to the toes and weakness in the lower extremity of the left leg. Diver recompressed to 60 fsw and treated on TT6 for DCS Type II. Complete relief noted 5 hrs 4 mins into treatment. No meds were used. 1 day loss of work and 28 days restricted from diving.

NMRI Diagnosis: Uncertain if decompression illness or not

Case 21*

This 29 year old male SEAL Diver made a SCUBA hookah dive from a Dry Deck Shelter (DDS) to 33 fsw for 4 hrs 38 mins. During drain down at 20 fsw diver noticed pain in his lower back. Drain depth was increased to 30 fsw for duration with no change noticed by diver. Upon surfacing a neuro exam was conducted with no other symptoms noted. Diver was recompressed to 60 fsw and treated on TT5 for DCS type I. Diver had complete relief 1 min into the treatment. No meds were used. No days loss of work and 2 days restricted from diving.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

Naval Safety Center Diagnosis: Type I Decompression Sickness

Case 22

This 22 year old male second class diver made a dive to perform a shaft wrap using MK 21. His dive profile was 25 fsw for 2 hrs 20 mins. While removing his gear on the surface the diver noticed that his right foot seemed to drop and he had decreased sensation over the dorsum of the right foot. Neuro exam was conducted with no other neurological symptoms noted. Diver recommpressed (*sic*) to 60 fsw and treated on a TT-6 for DCS type II. Diver had some relief 25 mins into the treatment but remained unchanged during the rest of the treatment. 1000 ML of Ringers Lactate administered IV. One day loss of work and 21 days restricted from diving.

NMRI Diagnosis: Uncertain if decompression illness or not

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 23

This 22 year old male second class diver made a recreational dive using open circuit scuba to 30 fsw for 45 mins. 2 hours after surfacing diver noticed intermittent tingling radiating from his right shoulder to his hand. Diver noticed right arm feeling numb the following morning and reported symptoms. Neuro exam revealed marked weakness to right arm and numbness of right shoulder to hand. Diver recompressed to 60 fsw and treated on TT 6 for DCS type II. No drugs were used. Diver had complete relief 3 hours and 14 mins into the treatment. 2 days loss of work and 30 days restructed (*sic*) from diving.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 24

This 38 year old male Master Diver made a ships husbandry dive using MK 21 to a depth of 22 fsw for 1 hr 31 mins. Diver surfaced with soreness in his right arm and feeling tired. Diver did not report these symptoms for two days. Neuro exam revealed right arm pain and soreness. Diver also was very fatigued. Diver recompressed to 60 fsw and treated on TT 6 for DCS Type II. No drugs were used. Diver had complete relief of symptoms 2 hrs 45 mins into the treatment. No days loss of work and 30 days restricted from diving.

NMRI Diagnosis: Uncertain if decompression illness or not

This 36 year old male civilian rigger diver made a ship's husbandry dive to 38 fsw for 1 hr 31 mins using MK 21. Upon surfacing the diver noticed the tip of his fifth digit on his left hand felt numb. Dive sup felt it may have been caused by a mechanical injury. After the diver had showered and warmed up he noticed the numbness had spread to his next finger. Diver transported to nearest chamber and recompressed to 60 fsw and treated on TT 6 for DCS Type II. No drugs were used. Diver had complete relief 23 mins into the treatment. No days loss of work and 14 days restricted from diving.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness or Arterial Gas Embolism)

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 26

This 23 year old male SEAL Diver made a SCUBA dive while conducting SDV diving operations. During the dive the diver reported that his wet suit was very tight and restrictive. Upon completion of his dive the diver noticed a patch of numbness on his left forearm. Neuro exam revealed no other symptoms. Diver was recompressed to 60 fsw and treated on TT 6 for possible DCS Type II. Diver had no relief of symptoms during the treatment. No days loss of work and 5 days restricted from diving.

NMRI Diagnosis: Definitely not decompression illness

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 27

This 24 year old male Navy SCUBA diver made a ship's husbandry dive to 45 fsw for 15 mins using open circuit SCUBA. Appox 1 and 1/2 hrs after surfacing diver complained of feeling "heavy" and tired. Diver was recompressed to 60 fsw and treated on TT 6 for DCS Type II. Diver had complete relief of symptoms 50 mins into the treatment. 4 Liters Ringer's Lactate IV was used. No days loss of work and 30 days restricted from diving.

NMRI Diagnosis: Uncertain if decompression illness or not

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 28

This 28 year old male First Class Diver made a ship's husbandry dive to 36 fsw for 1 hr 53 mins using MK -20. Approx 1 hr after surfacing diver developed a headache and fatigue. The next morning he reported his symptoms to the Diving Supervisor. Diver was recompressed to 60 fsw and treated on TT 6 for DCS type II. Diver had complete relief of symptoms 33 mins into the treatment. No drugs were used. No days loss of work ond 30 days restricted from diving.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

This 30 year old male USMC combat diver made a compass/navigation dive to 27 fsw for 1 hr 13 mins using open circuit SCUBA. Approx 30 mins after surfacing diver complained of tingling in his right forearm. Diver recompressed to 60 fsw and treated on TT6 for DCS type II. Diver had complete relief of symptoms 1 hr 54 mins into the treatment. No drugs were used. No days loss of work and 28 days restricted from diving.

NMRI Diagnosis: Uncertain if decompression illness or not Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 30

This 22 year old female student officer made a training dive using SCUBA. The dive was to 29 fsw for :29 min No D. Approx. upon surfacing she complained of a mild headache. A neuro was conducted by a DMO and no deficits were found. Oral motran (sic) was given at this time. Approx. one hour later she reported headache worsening, nauseous (sic), lightheaded and "not feeling right". A second neuro showed: pronounced weakness in left side of neck muscles, weakness in left extremities and degraded neural responses in left foot. Compressed to 165 fsw with 1st relief at 04: After leaving the surface. A TT 6A was completed with one extension at 60 fsw. An I.V. of Ringers Lactate was administered during treatment. The following AM left leg weakness was found on follow up neuro. A TT 6 was conducted with 1st relief at 1:29. A neuro on the following day still showed left leg weakness. A TT 6 was administered with complete relief at :28. Three days loss of work, 28 days restricted from diving, final diagnosis AGE. NMRI Diagnosis: Definitely decompression illness (Decompression Sickness or Arterial Gas

Embolism)

Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 31

This 20 year old male basic combat swimmer made an SDV training dive using SCUBA. The dive was to 40 fsw for 2 hrs and :01 min. Instructors observed him having difficulty maintaining depth control of the SDV. Upon surfacing the diver was unresponsive and aluggish (sic). The diving med tech on scene conducted a neuro and found upper extremity weakness. He was placed on surface 02 and transported to the chamber. He was recommpressed (sic) to 165 fsw where another neuro by a DMO showed upper extremity weakness, loss of coordination, proprioception and positive Babinski of right foot. Complete relief was obtained within :30 mins at 165 fsw. He was treated on a TT 6AM (sic), no meds were used, one day loss of work, 60 days restricted from diving, final diagnosis AGE.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness or Arterial Gas Embolism)

This 20 year old male Navy diving student made a training dive using open circuit SCUBA. The dive was to 24 fsw for :40 mins no "D". He surfaced complaining of feeling light headed. A neurological exam revealed no deficits. Approx. :30 minutes later he complained of pain in the left elbow. A neurological exam again showed no neurological deficits and the diver returned to the dive station. Approx. :20 to :30 the diver complained of tingling in left hand and fingers. Again the neuro was normal. AGE could not be ruled out so the diver was compressed to 165 fsw and treated on a TT 6A. Relief was obtained at :05 mins at treatment depth. No meds were used, no days loss of work and 28 days restricted from diving. Final diagnosis was AGE by DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 33

This 24 year old male UDT diver made a SPEC WAR ops dive using open circuit SCUBA. The dive was to 36 fsw for 50 minutes no "D" two minutes after surfacing this divers buddy experienced nausea and vomiting. A rapid upward excursion of approximately 12 fsw occurred during submarine lock in lock outs. A diagnosis of AGE could not be ruled out on the other diver so both divers were compressed to 165 fsw. 6 minutes after reaching treatment depth the afflicted diver was asymptomatic and neuros were normal in both divers. The divers air was obtained from the submarine's hp flasks and contamination could not be ruled out. An air sample was never recovered due to a mix up of SCUBA bottle during the transport of divers. No drugs were used one day loss of work and 30 days restricted from diving. Final diagnosis AGE by DMO.

NMRI Diagnosis: Definitely not decompression illness Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 34

This 21 year old UDT diver made a SPEC WAR ops dive using open circuit SCUBA. The dive was to 36 fsw for 50 minutes. Two minutes after surfacing this diver experienced nausea and vomiting. A rapid upward excursion of approximately 12 fsw occurred during submarine lock in lock out. A diagonosis of AGE could not be ruled out. This diver was compressed on to (sic) 165 fsw where he had complete relief after 6 minutes at treatment depth. A neuro was conducted at depth and was reported as normal. The divers (sic) air was obtained from the submarine's HP flasks and contamination could not be ruled out. An air sample was never recovered due to a mix up of SCUBA bottles during the transport of the divers. No drugs were used, one day loss of work, 30 days restricted from diving, final diagnosis AGE by DMO.

NMRI Diagnosis: Uncertain if decompression illness or not Naval Safety Center Diagnosis: Arterial Gas Embolism

This 18 year old male 2nd class diver made a searching dive using MK XXI air. The dive was to 26 fsw for 61 min. no "D". The members head being too small for the liner the oral nasal mask did not make a proper seal. This caused the mask to free flow. From 21 fsw to the surface this happened 3 times. Approx. 20 minutes after surfacing the diver noticed that he was feeling unusually tired he did not report this. 45 mins later he was found asleep. Upon being awakened it was noted that he had poor balance. A neuro produced a poor Romberg and patient was deteriorating rapidly. Compression to 165 commenced and complete relief was obtained at 100 fsw. A TT, (*sic*) 6A was completed. No meds were used, 3 days loss of work, 30 days restricted from diving, final diagnosis was AGE by DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness or Arterial Gas Embolism)

Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 36*

This 27 year old male 2nd class diver made a ships husbandry dive using ESDS air. The dive was to was to (*sic*) 37 fsw for 2 hours and 56 minutes no "D". Approx. one hour after surfacing the diver reported fullness around the neck, slight discomfort under sternum and tingling in left thoracic area. Initial neuro was normal. 25 minutes after the initial symptoms the diver reported pain in the left wrist. Another neuro revealed weakness in left leg. He was compressed to 60 fsw for treatment where the diver failed a Rhomberg Test. He was taken of (*sic*) O2 and compressed to 165 fsw (*sic*) All symptoms resolved at treatment depth. The Diver was treated on a TT 6A, and I.V. of Ringers Lactate was given. One days loss of work, 6 weeks restricted from diving, final diagnosis AGE by DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 37

This 23 year old 2nd class diver made a ships husbandry dive using MK 21 air. The dive was to 25 fsw for :44 min. no "D". Approx. 20 minutes after surfacing this diver experienced weakness, dizziness and fatigue. The dive was uneventful except he stated that his hat free floowed (*sic*) on two occasions during the dive. A neuro showed unsteady gait, Romberg Test was abnormal and a loss of feeling on portions of his back. A chest X-ray disclosed mediastinal emphysema. The diver was compressed to 60 fsw on O2 and had first relief after two minutes at 60 fsw with complete relief after the 2nd O2 period. A TT 6 was completed with one extension at 60 fsw. No meds were used, no days loss of work and 28 days restricted from diving. Final diagnosis possible AGE caused by improper fit of MK 21 diving helmet.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness or Arterial Gas Embolism)

This 25 year old male 1st class diver made a salvage dive using MK 21 air. The dive was to 33 fsw for 2:14 no "D". During the last three minutes of the dive this diver reported sudden onset of nausea, vertigo and headache. He vomited three times on the bottom. After several vents the diver was surface (*sic*). During the up and over phase the diver had trouble holding the ladder. A neuro conducted by the MDV confirmed weakness in shoulder and elbow muscle groups on left side. He was transported on O2 to chamber. Compressed to 165 fsw with first relief at 2 minutes on the bottom. Relief of all symptoms at 29 minutes. A TT 6A was completed. No meds were used, no days loss of work, 28 days restricted from diving. Final diagnosis hypercapnia and AGE.

NMRI Diagnosis: Uncertain if decompression illness or not Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 39*

This 28 year old male 1st class diver made a ships husbandry dive using ESDS (AGA) on air. The dive was to 34 fsw for 3:11 no "D". Approx 15 to 20 minutes after the dive this diver noticed a loss of hearing in his left ear. No other neurological deficits were noted. Diver was compressed to 165 fsw. EDU was notified of mishap and concurred with treatment. First relief was noted at 1:45 into the treatment at 60 fsw on O2. A TT 6A was completed with no extensions. The diver surfaced with substantial relief. No meds were used, one day loss of work and 30 days restricted from diving. Final diagnosis was AGE by DMO.

NMRI Diagnosis: Uncertain if decompression illness or not Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 40

This 24 year old female diving officer made searching dive using open circuit SCUBA. The dive was to 40 fsw for 3 minutes. While at depth this diver reported vertigo. The dive was aborted. On the surface she complained of dizziness and severe headache. A neuro revealed neurological deficits heel to toe walk. The diver was compressed to 165 fsw. 1st relief was noted upon reaching treatment depth. Complete relief was noted 2 minutes into the 2nd O2 period at 60 fsw. (*sic*) and I.V. of normal saline and I.V. decadron was given. No days loss of work, 28 days restricted from diving, final diagnosis AGE by MDV.

NMRI Diagnosis: Definitely not decompression illness Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 41

This 20 year old 2nd class diver made a salvage dive using open circuit SCUBA. The dive was to 35 fsw for :51 BT. This diver was not acting normal after the dive. A neuro was conducted. He could not pass the Romberg portion of the exam. 11 minutes after surfacing he was compressed to 165 fsw. :13 minutes into the treatment complete relief was obtained. A TT 6A was completed, no meds were used, one day loss of work and 28 days restricted from diving. Final diagnosis was AGE by DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness or Arterial Gas Embolism)

This 3M (*sic*) year old 2nd class diver made a salvage dive using open circuit SCUBA. The dive was to 35 fsw for :52 BT. Approx. eight minutes after surfacing this diver reported muscle tightness (mechanical injury) in the right arm. He was swinging a hammer at depth. A neuro was performed by a DMO. AGE could not be ruled out for his diving buddy showed neurological deficit in his neuro exam. This diver was compressed to 165 fsw 2 and one half hours after surfacing. A TT 6A was completed with full relief noted :10 into the treatment. No meds were used, one day loss of work and 28 days restricted from diving. Final diagnosis was AGE by DMO.

NMRI Diagnosis: Uncertain if decompression illness or not Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 43

This 21 year old male Navy EOD assistant diver made an EOD dive using open circuit SCUBA. The dive was to 50 fsw for :40 BT no "D". This diver make (*sic*) 3 rapid ascents to the surface from 40 fsw to retrieve buoyant objects. He exceeded 60 fpm on ascent passing his own bubbles. Blood was in his face mask upon surfacing. Symptoms of numbness both hands and feet were noted five hours post dive but not reported until 15 hours after surfacing. He was compressed to 60 fsw on O2, 1st relief was obtained upon reaching treatment depth, complete relief at 1:00 into the treatment. TT 6 with 1 ext at 60 fsw was completed. No meds were used, one day loss of work and 28 days restricted from diving. Final diagnosis was DCS II by DMO. Note: reported as DCS II. The dive profile and type of symptoms strongly suggest AGE.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

Naval Safety Center Diagnosis: Type II Decompression Sickness

Case 44

This 37 year old male Navy Sat Diving Officer made a ships husbandry dive using MK 21 on air. The dive was to 30 fsw for :25. After the dive the Diving Supervisor, which was under instruction, performed a training neuro on this diver which revealed a neurological deficit. A lack of communication was aparent (sic). The deficit was thought to be manufactured by the diver for the training neuro. The nest (sic) day he was evaluated for the deficit. It consisted of personality changes. Lack of coordination and a positive Babinski sign. He was compressed to 60 fsw on O2. Complete relief was noted :19 into the treatment. An I.V. of NACL was used, 2 days loss of work, 35 days restricted from diving. Final diagnosis was AGE by DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness or Arterial Gas Embolism)

Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 45

This 21 year old male Navy diving student made a training dive using Mark 21 on air. The dive was to 29 fsw for :22 BT. :08 after surfacing this diver complained of a headache. A neuro was normal. The diver stated that his headache was worsening. He was compressed to 165 fsw. He reported complete relief on descent at 18 fsw. No meds were used, no days loss of work and 28 days restricted from diving. Final diagnosis was AGE by DMO.

NMRI Diagnosis: Uncertain if decompression illness or not Naval Safety Center Diagnosis: Arterial Gas Embolism

This 23 year old male SPECWAR Diver made a lock in lock out dive using open circuit SCUBA. The dive was to 29 fsw for 1:14. The diver used a lift bag to float an outboard engine to the surface. Upon surfacing he experienced dizziness. He made a descent for lock in. The neuro was normal except for dizziness. He was compressed to 165 fsw. First relief was noted :02 into the treatment at 60 fsw. Complete relief was noted :10 into the treatment. A TT 6A was completed. An I.V. of Ringers D5W was given. 3 days loss of work, 42 days restricted from diving. Final diagnosis was AGE by DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness or Arterial Gas Embolism)

Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 47

This 23 year old male 2nd class diver made a ships husbandry dive using ESDS on air. The dive was to 35 fsw for 1:55. This diver (red) noted that he was not feeling well on the bottom and turned over his work to his dive buddy (yellow). Red divers (*sic*) condition did not improve. The supervisor recovered both divers. Once on the surface yellow diver reported the same feeling earlier but it cleared. Red diver on the surface complained of nausea and light-headedness. At the chamber a neuro showed short-term memory loss, difficulty in talking and nystagmus. He was compressed to 165 fsw. First relief was noted upon reaching treatment depth and complete relief at 2:14 into the treatment. A TT 6A was completed. No meds were used, no day loss of work and 30 day restricted from diving. Final diagnosis was AGE by DMO. Cranial nerve deficite (*sic*)

NMRI Diagnosis: Uncertain if decompression illness or not Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 48

This 23 year old male EOD assistant made an EOD OP dive using open circuit SCUBA. The dive was to 45 fsw for :23 BT. While on the bottom this diver experienced a headache and surface (*sic*). On the surface he had extreme weakness and had difficulty remaining alert. He was examined by DMO and MDV. A diagnosis of AGE was made. He was transported to the chamber and compressed to 165 fsw. Complete relief was noted at 130 fsw on descent. At (*sic*) TT 6A was completed. No meds were used, no days loss of work and 42 days restricted from diving.

NMRI Diagnosis: Definitely decompression illness (Arterial Gas Embolism)

This 40 year old male civilian diver made an inspection dive using open circuit SCUBA and a dry suit. The dive was to 47 fsw for 5M mins. 20 mins after surfacing this diver complained of dizziness and nausea. He was diagnosed as having shallow water AGE by DMO. Recompressed to 60 fsw. First relief occured (sic) 4 mins into the treatment followed by complete relief 5 hours and 36 mins into the treatment. A TT6 was completed. No meds were used. No days loss of work and 30 days restricted from diving.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness or Arterial Gas Embolism)

Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 50

This 23 year old male second class diver made a SSBN security swim using surface tended open circuit SCUBA. Diver's tending line snagged on rudder and as he attempted to unsnag himself he made an uncontrollable ascent (8 to 10 fsw) to the surface. Dive profile was 40 fsw for 20 mins. Approximately 3 mins after surfacing diver complained of numbness in lips and facial area. While transiting to the chamber diver reported numbness and loss of strength to left arm and shoulder, dizziness, and blurred vision. Seconds prior to entering chamber diver became unconsciousness (sic). Recompressed to 165 fsw and treated on TT6A for AGE. Complete relief noted 5 M mins into the treatment. 1000ML 2% sodium chloride IV used. Three days later this same diver reported feeling fatigued, headache, back pain and left elbow pain. Neuro exam conducted revealing pain in back of head, pain middle left side of back, poor coordination, tingling left hand, and slurred speech. Diver recompressed to 60 fsw on a TT6. Complete relief noted 40 mins into the treatment. No meds were used. No days loss of work and 2 days restricted from diving.

NMRI Diagnosis: Definitely decompression illness (Arterial Gas Embolism)

Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 51

This 38 year old male HeO2 diving officer was assisting during a recovery of a DSRV using open circuit SCUBA. His dive profile was 42 fsw for 20 mins. 3 mins after surfacing diver became unconscious and had a grand mal seizure. Seizure lasted approximately 1 min and diver regained consciousness about 15 mins later. Diver medivaced to chamber by helo. Neuro exam conducted with no neurological symptoms noted. Diver recompressed to 60 fsw on TT6 for AGE. Ringers lactated (sic) used during treatment. No days loss of work and 42 days restricted from diving.

NMRI Diagnosis: Definitely decompression illness (Arterial Gas Embolism)

This 22 year old male second class diver was patching a wreck using MK 21. His dive profile was 52 fsw for 58 mins. 30 mins after surfacing diver complained of pain in his lower back and stated that he did not feel well. Neuroexam was conducted and a loss of hand to eye coordination and loss of balance was noted. He was recommpressed (*sic*) to 165 fsw on a TT6A for AGE. Neuro exam at 165 fsw noted short term memory loss, weakness in the left arm, and drooping facial muscles. Complete relief noted 23 mins into the treatment. No meds were used. Upon surfacing neuro exam revealed roving left eye. Diver recompressed to 60 fsw and treated on TT6. Complete relief noted 19 mins. into the treatment. No meds used. 1 day loss of work and 28 days restricted from diving.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness or Arterial Gas Embolism)

Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 53

This 21 year old male dive student was conducting underwater sonar operations using a hand held Dukane sonar while diving SCUBA. His dive profile was to 22 fsw for 25 mins. Upon handing the sonar to his dive partner he noticed tingling in his 1st and 2nd fingers on his right hand. Two minutes later he aborted the dive and reported his symptoms to the Diving Supervisor. Diver was transported to the chamber and examined by DMO where he was diagnosed as having fatigue/altered (*sic*) sensation to his right hand due to holding the sonar. DO elected to treat the diver for AGE due to neurological symptom being present. Attempted to recompress diver to 165 fsw but diver had problems clearing and descent was stopped at 71 fsw and diver brought back up to 60 fsw and treated on TT-6. Diver had complete relief of all symptoms at 39 fsw on descent. No meds were used. One day loss of work and 28 days restricted from diving.

NMRI Diagnosis: Definitely not decompression illness Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 54

This 24 year old male first class diver made a dive to seal a cofferdam which had developed a leak using MK 20. As the diver was returning to the cofferdam, it blew out a two foot section of the rubber releasing a large air pocket which hit the diver forcing him to the surface. His dive profile was to 28 fsw for 37 mins. Upon surfacing the diver was semi-conscious. A neuro exam was conducted which revealed tingling and loss of sensation in the right upper extremity from the shoulder down phrough (sic) the hand and the left wrist down through the hand. The diver also complained of nausea. Diver was recompressed to 60 fsw and treated on a TT-6 for AGE. Diver had complete relief 22 mins into the treatment. 1000 ml Ringers Lactate and 10 mg Dexamethasone given IV. 1 day loss of work and 28 days restricted from diving.

NMRI Diagnosis: Definitely decompression illness (Arterial Gas Embolism)

This 26 year old male first class diver made a bottom search dive using MK-20 to 35 fsw for 43 mins. 40 mins. after surfacing the diver complained of pain in his left elbow. A neuro exam was conducted and weakness of his left arm and shoulder were discovered as well. Diver was put on O2 and transported to nearest recompression chamber. Diver had complete relief of symptoms enroute to the chamber. Diver recompressed to 60 fsw and treated on TT 6 for AGE. No meds were used. 1 day loss of work and 28 days restricted from diving.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 56

This 22 year old male second class diver made a dive to inspect a shaft system using SCUBA to a depth of 23 fsw for 44 mins. Approx 26 hours after surfacing diver complained of pain and weakness of left foot and tightness across right chest area. Diver recompressed to 60 fsw and treated on TT 6 for AGE. Diver had complete relief five minutes into the treatment. No meds were used. No days loss of work and 28 days restricted from diving.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 57

This 26 year old male EOD diver made an open circuit SCUBA dive to 37 fsw for 20 mins using a drysuit. During the dive the diver experienced difficulty controlling the buoyancy of his drysuit. Approx 1 hr after the diver the diver complained of a dull pain in his neck apound (sic) the area of the spinal cord. A neuro exam was conducted with no other symptoms noted. Diver recompressed to 60 fsw and treated on a TT6 for AGE. No meds were used. No days loss of work and 42 days restricted from diving.

NMRI Diagnosis: Uncertain if decompression illness or not Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 58

This 22 year old male EOD diver made an open circuit SCUBA dive to 40 fsw for 20 mins. During the dive the diver lost his weight belt and was unable to control his ascent to the surface. Upon surfacing the diver felt disorientated (*sic*). Neuro exam revealed weakness of right arm and leg. Diver recompressed to 60 fsw and treated on TT 6 for AGE. No drugs were used. Diver had complete relief 51 mins into the treatment. No days loss of work and 14 days restricted from diving.

NMRI Diagnosis: Definitely decompression illness (Arterial Gas Embolism)

This 34 year old male Navy SPECWAR SOT tech made a SPECWAR dive using open circuit SCUBA. The dive was to 25 fsw for :40. Forty minutes after surfacing he complained of pain in the left hand. This may have been present during the dive and probably due to a grip maintained during the dive. The exam was normal and told to return if symptoms worsened. Later on that night he noticed paresthesia in the hand, a funny feeling in his left elbow, shoulder and lower back. He called in and was told to report ASAP. The exam this time was positive for neurological deficits. He was compressed to 165 fsw. First relief was noted :11 into the treatment and complete relief :16 later. A TT 6A with extensions at 60 and 30 fsw was completed. No meds were used, 2 days loss of work and 14 days restricted from diving. Final diagnosis was AGE by DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)

Naval Safety Center Diagnosis: Arterial Gas Embolism

Case 60

This 19 year old male second class Navy diver made a ships husbandry dive using Mark 20. The dive was to 21 fsw for one hour and three minutes, no "D". The next AM while working out at the gym this diver noticed slight numbness and tingling in his right hand. He stopped his exercise and returned to the dive locker. His symptoms exacerbated during transit including numbness and loss of strength in the lower right extremity and a decrease in peripheral vision. The neuro at the chamber revealed six significant deficits. He was compressed to 60 fsw on O2. First relief was noted 8 minutes in to the treatment and complete relief during the second O2 period. A treatment table 6 was completed. No meds were used, no days loss of work and 28 days restricted from diving. Finial (sic) diagnosis was DCS Type II by the DMO.

NMRI Diagnosis: Definitely decompression illness (Decompression Sickness)